

3rd International Conference and Exhibition on **COSMETOLOGY & Trichology**

July 21-23, 2014 Hampton Inn Tropicana, Las Vegas, USA

Autogenous adipose tissue with ADSC for mammaplasty and renovation

D S Johnson Chang

YAyo Aesthetic, Canada

Beautiful face according to: Culture, generation, gender, age and Darwin's face average: Balance, symmetry and harmonize.

Nose special points: Lower and wide of nasal bridge, shorter, fatty, lower nasal tip, thicknesses and less nasal cartilage.

Face special points: Empty of temporal area, lower and unsmooth of forehead, higher zygomatic area, empty of around eyes and lower or shorter chin.

Methods: Autogenous adipose tissue with ADSC augmentation: 10 ml to 105 ml to area of forehead, chin, temporal area, peri orbital area, smiling m., Cheek, nasal base, nasal labial, peri oral area, ear drop, lips, rhino area.

Size of dimension lines: 22 mm, 26mm, 28mm, 30mm, 32mm, 36mm.

Laser liposuction: LAS - 30B, duration 100 msec, frequency 8 Hz, aim beam 10, output power 5-8 w Rhinoplasty, blepharoplasty

Conclusion: The compose some aesthetic methods for face rejuvenation that can enhance coordinating and sense of beauty. The Rhinoplasty complication with un- smooth or shadow of implant that can use Autogenous adipose tissue be adjustment filler. Autogenous adipose tissue is a filler and also permanent filler. Laser liposuction can de-fat double chin and jaw line and also can lift of whole face. Dimension line can be a construction function that become a space to restore filler or Autogenous adipose tissue, it will be absorptive in 8 month but Autogenous adipose tissue will regrow on site in 2 month after transplantation.

Biography

D S Johnson Chang has 26 years of experience in autogenus adipose tissue transplantation. He had done his plastic surgery training course for 8 years in University Hospital after graduation at Medical School in Taiwan. And have done his PhD degree of stem cell treatment in PU on 2003.

johnsonchangmd@icloud.com