

A Comparative Biochemical Analysis of Photosynthetic Efficiency in C3 and C4 Plants

Lewis Jane *

Department of Agricultural Sciences, Stanford University, California, United States of America

DESCRIPTION

Photosynthesis is a critical process for plant life and global carbon cycling, primarily occurring in two pathways: C3 and C4 photosynthesis. This article describes the to provide a comparative biochemical analysis of the photosynthetic efficiency of C3 and C4 plants, highlighting the underlying mechanisms and adaptations that enable each pathway to optimize carbon fixation under varying environmental conditions [1,2].

Photosynthesis is the biological process by which light energy is converted into chemical energy, primarily through the conversion of Carbon Dioxide (CO₂) and water into glucose and oxygen. The two primary pathways for photosynthesis, C3 and C4, exhibit distinct biochemical and physiological characteristics, influencing their efficiency under different environmental conditions. C3 plants, such as wheat and rice, use ribulose biphosphate carboxylase/oxygenase (Rubisco) for CO₂ fixation, while C4 plants, such as maize and sugarcane, utilize a two-step process involving Phosphoenolpyruvate Carboxylase (PEPC) and a specialized bundle sheath cell architecture to concentrate CO₂. This communication analyzes the biochemical differences between these two pathways and their implications for photosynthetic efficiency [3,4].

Biochemical mechanisms of photosynthesis

C3 photosynthesis: C3 photosynthesis occurs in three primary stages: Light reactions, the Calvin cycle and regeneration of Ribulose Biphosphate (RuBP). During the light-dependent reactions, chlorophyll pigments absorb light energy, driving the synthesis of Adenosine Triphosphate (ATP) and Nicotinamide Adenine Dinucleotide Phosphate (NADPH). In the Calvin cycle, Rubisco catalyzes the reaction of CO₂ with RuBP, resulting in the formation of 3-Phosphoglycerate (3-PGA), which is subsequently converted into glucose through a series of enzymatic reactions [5,6].

C4 photosynthesis: In contrast, C4 photosynthesis features a more complex pathway to enhance CO₂ fixation and reduce

photorespiration. The process begins with the enzyme PEPC, which captures CO₂ to form Oxaloacetate (OAA) in mesophyll cells. OAA is then converted into malate or aspartate, which is transported to bundle sheath cells, where it is decarboxylated to release CO₂ for the Calvin cycle. This compartmentalization allows C4 plants to maintain a higher concentration of CO₂ around Rubisco, enhancing photosynthetic efficiency, especially under high light intensity, elevated temperatures and arid conditions [7].

Comparative photosynthetic efficiency

Efficiency under optimal conditions: Under optimal conditions, C4 plants demonstrate significantly higher photosynthetic efficiency than C3 plants. Studies have shown that C4 photosynthesis can achieve up to 30% greater photosynthetic rates due to its ability to concentrate CO₂. This increased efficiency is particularly beneficial in environments with high temperatures and intense sunlight, where photorespiration in C3 plants can lead to substantial carbon loss [8,9].

Stress responses: C3 plants are generally more efficient under cooler and wetter conditions, while C4 plants are adapted to hotter and drier climates. C3 plants exhibit a higher carboxylation efficiency under low CO₂ concentrations, making them well-suited for shaded or suboptimal environments. In contrast, C4 plants maintain higher rates of photosynthesis and lower rates of photorespiration in high light and temperature conditions [10].

CONCLUSION

In summary, the comparative biochemical analysis of C3 and C4 photosynthetic pathways reveals distinct mechanisms that confer varying efficiencies under specific environmental conditions. C4 plants possess adaptations that enhance their performance in high light and temperature scenarios, while C3 plants exhibit advantages in cooler, shaded environments. Understanding these differences is essential for improving agricultural productivity and

Correspondence to: Lewis Jane, Department of Agricultural Sciences, Stanford University, California, United States of America, E-mail: jane_lewis@gmail.com

Received: 16-Mar-2025, Manuscript No. JPBP-24-34913; **Editor assigned:** 19-Mar-2025, PreQC No. JPBP-24-34913 (PQ); **Reviewed:** 02-Apr-2025, QC No. JPBP-24-34913; **Revised:** 09-Apr-2025, Manuscript No. JPBP-24-34913 (R); **Published:** 16-Apr-2025, DOI: 10.35248/2329-9029.25.13.353

Citation: Jane L (2025). A Comparative Biochemical Analysis of Photosynthetic Efficiency in C3 and C4 Plants. J Plant Biochem Physiol.13:353.

Copyright: © 2025 Jane L. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

developing strategies to enhance crop resilience in the face of climate change.

REFERENCES

1. Abdelgawad H, Hassan YM, Alotaibi MO, Mohammed AE, Saleh AM. C3 and C4 plant systems respond differently to the concurrent challenges of mercuric oxide nanoparticles and future climate CO₂. *Sci Total Environ.* 2020;749:142356.
2. Andersen JR, Lübberstedt T. Functional markers in plants. *Trends Plant Sci.* 2003;8(11):554-560.
3. Almeselmani M, Deshmukh PS, Sairam RK, Kushwaha SR, Singh TP. Protective role of antioxidant enzymes under high temperature stress. *Plant Sci.* 2006;171(3):382-388.
4. Alton P, Bodin P. A comparative study of a multilayer and a productivity (light-use) efficiency land-surface model over different temporal scales. *Agric For Meteorol.* 2010;150(2):182-195.
5. Elgawad H, de VD, Zinta G, Domagalska MA, Beemster GT, Asard H. Grassland species differentially regulate proline concentrations under future climate conditions: An integrated biochemical and modelling approach. *New Phytol.* 2015;208(2):354-369.
6. Singh RK, Muthamilarasan M, Prasad M. Biotechnological approaches to dissect climate-resilient traits in millets and their application in crop improvement. *J Biotechnol.* 2021;327:64-73.
7. Alexandridis TK, Ovakoglou G, Clevers JG. Relationship between MODIS EVI and LAI across time and space. *Geocarto Int.* 2020;35(13):1385-1399.
8. Seki M, Umezawa T, Urano K, Shinozaki K. Regulatory metabolic networks in drought stress responses. *Curr Opin Plant Biol.* 2007;10(3):296-302.
9. Atchudan R, Edison TN, Shanmugam M, Perumal S, Somanathan T, Lee YR. Sustainable synthesis of carbon quantum dots from banana peel waste using hydrothermal process for *in vivo* bioimaging. *Physica E Low Dimens. Syst. Nanostruct.* 2021;126:114417.
10. Akdemir KC, Chin L. HiCPlotter integrates genomic data with interaction matrices. *Genome Biol.* 2015;16:1-8.