

Cardiac Specific Knockout of P65 Mice Resist to Cardiac Ischemia-Reperfusion Injury

Xiu Zhang*

Department Obstetrics and Gynecology, University of Utah School of Medicine, Utah, USA




Figure 1: (A) Hearts from mice exposed to left anterior descending coronary artery permanent ligation ischemia for 7 days were harvested and then stained with tri-chrome staining. Fibrosis of infarct area was stained to blue and viable tissue was stained to red. (B) Myocardial Infarction fraction was determined using ImagePro analysis software. (XiuQuan Zhang, MD/PhD, University of Utah School of Medicine).

*Corresponding author: Xiu Zhang, Department Obstetrics and Gynecology, University of Utah School of Medicine, 50 N Medical Dr. Salt Lake City, Utah 84132, USA, Tel: 801-585-3117; E-mail: iuquan.zhang@hsc.utah.edu

Received October 31, 2014; Accepted October 31, 2014; Published November 02, 2014

Citation: Zhang X (2014) Cardiac Specific Knockout of P65 Mice Resist to Cardiac Ischemia-Reperfusion Injury. Adv Genet Eng 3: i101. doi:[10.4172/2169-0111.1000i101](https://doi.org/10.4172/2169-0111.1000i101)

Copyright: © 2014 Zhang X. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.