

Evaluation of *Stylosanthes* Species and their Accessions for Herbage Yield in Two Agro-Ecologies of Benishangul-Gumuz Region, Western Ethiopia

Mulisa Faji*, Alemeyehu Abebe, Kedija Ahmed, Workene Tezera

Department of Agricultural Research, Ethiopian Institute of Agricultural Research, Addis Ababa, Ethiopia

ABSTRACT

Three *Stylosanthes hamata* accessions and one accession of *S. scabra* were evaluated for their agronomical performance under two environmental conditions of Benishangul-Gumuz region state, Western Ethiopia. The experiment was carried out at Kamash and Assosa Forage research station of Assosa agricultural research center and were purposively selected to represent lowland and mid-altitude agro-ecologies, respectively. The experiment was conducted in a randomized complete block design with three replications. The general linear model procedures of SAS and least significance difference for data analysis and mean separation were employed, respectively. Environment was significantly ($P < 0.001$) affecting dry matter yield, number of tillers, number of branches per plants and leaf to stem ratio. The highest forage dry matter yield was obtained at Assosa. The results of the combined analysis revealed that plant height at forage harvesting ($P < 0.05$), number of tillers ($P < 0.01$) and leaf to stem ratio ($P < 0.01$) were significantly influenced by genotype. Taller plant height and higher leaf to stem ratio were obtained from *Stylosanthes scabra* 441. Leaf to stem ratio was significantly ($P < 0.01$) different among genotypes of Assosa and the highest leaf to stem ratio was recorded for *Stylosanthes scabra* 441. The interaction effect of location and genotype was nonsignificant ($P > 0.05$) for all measured parameters and this indicated consistency in the performance of the genotypes across the environment and this leads to conclude no need for assessing performance to identify *Stylosanthes* genotypes with stable and superior yield across the environments.

Keywords: *Stylosanthes*; Dry matter yield; Environment; Genotype; Leaf to stem ratio

INTRODUCTION

Inadequate supply and quality of forage, particularly during the dry season, is a major constraint to livestock production in Sub-Saharan Africa [1]. Poor management of the available feeds, seasonal variability in weather and climate change contributes to the high fluctuation of forage quality and quantity between seasons and years. To improve livestock production, there is a need to improve both the quantity and quality of available feed through the use of alternative forage crops like *Stylosanthes*. *Stylosanthes* is regarded as the most economic and significant pasture and forage legume in the tropical regions. Consequently, as an attempt to improve livestock nutrition, it was introduced in West Africa in the 1960's after several new improved cultivars were developed in Australia [2].

Stylosanthes legume has been the forage of interest in Africa for pasture improvement, particularly West Africa. This is based on the merits of the genus, which include high yield of protein per hectare [5]. According to [6], the genus *Stylosanthes* has provided ample germplasm for a wide variety of agro-ecological situations in the tropics. *Stylosanthes* has fitted successfully to the dry land of African agriculture, particularly because of its drought-tolerant characteristics [3]. Therefore, based on the merits of *Stylosanthes* genus in improving feed quality and soil fertility, an evaluation of some major *Stylosanthes* species plays a paramount role to alleviate the feed problems in quality and quantity. Consequently, this study was conducted with the aim of determining the adaptability and agronomic potential of four *Stylosanthes* for age genotypes under the two agro-ecological zones of Benishangul-Gumuz regional state, Western Ethiopia.

Correspondence to: Mulisa Faji, Department of Agricultural Research, Ethiopian Institute of Agricultural Research, Addis Ababa, Ethiopia; Email: tugbagungor@comu.edu.tr

Received: 15-Apr-2020, Manuscript No. JBFBP-24-3888; **Editor assigned:** 20-Apr-2020, PreQC No. JBFBP-24-3888 (PQ); **Reviewed:** 04-May-2020, QC No. JBFBP-24-3888; **Revised:** 22-May-2024, Manuscript No. JBFBP-24-3888 (R); **Published:** 19-Jun-2024, DOI: 10.35248/2593-9173.24.15.173.

Citation: Faji M, Abebe A, Ahmed K, Tezera W (2024) Evaluation of *Stylosanthes* Species and their Accessions for Herbage Yield in Two Agro-Ecologies of Benishangul-Gumuz Region, Western Ethiopia. J Agri Sci Food Res. 15:173.

Copyright: © 2024 Faji M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

MATERIALS AND METHODS

Study area

The trial was conducted under field conditions at Assosa and Kamash forage research stations of Assosa agricultural research center during the main cropping season.

The test locations represent the mid-and low altitude areas ranging in altitude from 1000 m.a.s.l to 1576 m.a.s.l. The farming system of the study area is Agro pastoral. Descriptions of the test environments are indicated in Table 1 [4].

Table 1: Description of the test environments for geographical position and physico-chemical properties of the soil.

| Parameters | Study sites | |
|--------------------------------|-------------|-----------|
| | Kamash | Assosa |
| Latitude | 09°30'N | 10°30'N |
| Longitude | 35°45'E | 034°20'E |
| Altitude (masl) | 1000-1350 | 1500-1576 |
| Annual rainfall (mm) | 1150 | 1316 |
| Daily minimum temperature (°C) | 25 | 16.75 |
| Daily maximum temperature (°C) | 30 | 27.9 |

Experimental treatment and design

The four genotypes of *Stylosanthes* for this research experiment were collected from ILRI. The genotypes were planted in a 3 m × 4 m plots using a Randomized Complete Block Design (RCBD) with three replications at the beginning of the main rainy season. Seed was sown 30 cm spacing between rows by drilling at 3 cm depth. The total experimental area was 13 m × 20.5 m (266.5 m²) with an individual plot size of 12 m² and spacing between plots and replications of 1.5 m and 2 m, respectively at each testing environment. The treatments were sown according to their recommended seeding rates: 2 kg ha⁻¹-10 kg ha⁻¹ and fertilizers were not applied [5].

Data collection

Data was collected on number of tillers, Plant Height at Harvesting (PHH), number of branches per plant and forage dry matter yield. Number of tillers, plant height at harvesting and number of branches per plant were taken for six plants randomly selected from each plot. Plant height at harvesting was measured using a steel tape from the ground level to the highest leaf. For determination of biomass yield, genotypes were harvested at the forage harvesting stage (50% blooming stage) in the laid quadrant which has 1 m² area. Weight of the total fresh biomass yield was recorded from each plot in the field and the estimated 500 g sample was taken from each plot to the laboratory. The sample taken from each plot was weighed to know their sample fresh weight and oven dried for 72 hours at a temperature of 65°C to determine dry matter yield [6].

Statistical analysis

Analysis of Variance (ANOVA) procedure of SAS General Linear Model (GLM) was used to compare treatment means.

LSD test at 5% significance will be used for comparison of means. The Pearson correlation analysis procedure of the SAS statistical package was applied to measure the strength of linear dependence between any two measured variables. The data was analyzed using the following model:

$$Y_{ijk} = \mu + G_i + E_j + GE_{ij} + B_k + e_{ijk}$$

Where, Y_{ijk} = Dependent variables

μ = Grand mean

G_i = Effect of genotype i

E_j = Effect of environment j

j = Assosa and Kamash

GE_{ij} = The interaction effect of genotype i and environment j

B_k = Effect of block k, and

e_{ijk} = Random error effect of genotype i, environment j, interaction effect of genotype i and environment j, and block k

RESULTS AND DISCUSSION

Environment and interaction effect on *Stylosanthes* genotype performance

Combined analysis of variance for measured agronomic traits of *Stylosanthes* genotypes tested over environment is presented in Table 2. The results of this study revealed that the environment was significantly ($P < 0.001$) influencing the forage dry matter yield, number of tillers, number of branches per plant and leaf to stem ratio of the tested *Stylosanthes* genotypes. Plant height at forage harvesting ($P < 0.05$), number of tillers ($P < 0.01$) and leaf to stem ratio ($P < 0.01$) were significantly affected by genotype.

Interaction of genotype and environment was not significantly ($P>0.05$) affecting the measured agronomic traits except leaf to stem ratio ($P<0.05$) and this indicated consistency in the performance of the genotypes across the environments and this leads to no need for assessing performance to identify *Stylosanthes* genotypes with stable and superior yield across the environments. Statistically, $G \times E$ interactions are detected as significantly different patterns of response among genotypes across environments, this will occur when the contributions (or level of expression) of the genes regulating the trait differ among environments [7].

The forage dry matter yield performance of the tested *Stylosanthes* genotypes was stable across the environment and this might be due to the interaction effect of genotype and environment that was nonsignificantly influencing the forage dry matter yield. In agreement with this study was reported that a major difference in genotype stability is due to the crossover interaction effect of genotype and environment.

Table 2: Combined analysis of variance for measured agronomic traits of four *Stylosanthes* genotypes tested across two locations/environments.

| Traits | Mean square | | $G \times E$ | Mean | CV |
|---------------------------|-------------|-------------|--------------|-------|-------|
| | Genotype | Environment | | | |
| Plant height (cm) | * | ns | ns | 42.54 | 19.37 |
| Forage DM yield (t/ha) | ns | ns | ns | 3.67 | 51.7 |
| Number of tiller | ** | *** | ns | 3.9 | 36.24 |
| No. of branches per plant | ns | *** | ns | 9.44 | 23.99 |
| Leaf to stem ratio | ** | *** | * | 0.61 | 18.65 |

Note: $G \times E$ =Interaction of genotype and environment; CV=Coefficient Variation; ns=non-significant; * = $P<0.05$; **= $P<0.01$; ***= $P<0.001$

Leaf to stem ratio

The mean leaf to stem ratio at forage harvesting of the four tested *Stylosanthes* genotypes under the two agro-ecological zones of Benishangul-Gumuz is presented in Table 3. The result of combined ($P<0.05$) and Assosa location ($P<0.01$) analysis indicated that the leaf to stem ratio was significantly different among the tested genotypes and *Stylosanthes scabra* 441 was given the highest value.

Mean leaf to stem ratios of the tested *Stylosanthes* genotypes were significantly ($P<0.001$) affected by location. The results indicated that the highest mean leaf to stem ratio at forage harvesting stage was recorded from Kamash.

Table 3: Mean leaf to stem ratio of four *Stylosanthes* genotypes tested across two locations/environments at forage harvesting stage.

| Genotypes | Location/Environments | | Combined analysis |
|--------------------------------|-----------------------|--------|-------------------|
| | Assosa | Kamash | |
| <i>Stylosanthes hamata</i> 167 | 0.46 | 0.69 | 0.58 |

According to the interaction is a result of changes in a cultivar's relative performance across environments due to differential responses of the genotypes to various edaphic, climatic and biotic factors. When genotypes perform consistently across locations, breeders are able to effectively evaluate the germplasm with a minimum cost in a few locations for the ultimate use of the resulting varieties across wider geographic areas. However, with high genotype by location interaction effects, genotypes selected for superior performance under one set of environmental conditions may perform poorly under different environmental conditions. Therefore, the development of cultivars or varieties, which can be adapted to a wide range of diversified environments, is the ultimate goal of plant breeders in a crop improvement program. Therefore, according to the finding of these authors, evaluation of yield, performance and adaptation patterns of *Stylosanthes* genotypes in multiple environments could't be important step in agronomic evaluation and selection of better adapted and high yielding species and varieties [8].

| | | | |
|----------------------------------|--------|-------------------|--------|
| <i>Stylosanthes hamata</i> 75 | 0.38 | 0.75 | 0.56 |
| <i>Stylosanthes hamata</i> 15876 | 0.42 | 0.73 | 0.57 |
| <i>Stylosanthes scabra</i> 441 | 0.69 | 0.76 | 0.72 |
| Mean | 0.49 | 0.73 ^a | 0.61 |
| CV | 25.35 | 14.79 | 29.73 |
| P-value | 0.0017 | 0.7315 | 0.0142 |

Plant height at forage harvesting

The mean plant height at forage harvesting of the four tested *Stylosanthes* genotypes is presented in Table 4. Mean plant height at forage harvesting was non-significantly ($P>0.05$) different across the testing environments. The results of the combined analysis indicated that the mean plant height at forage harvesting was significantly ($P<0.05$) different among the tested genotypes. Among the tested *Stylosanthes* genotypes, the highest mean plant

height at forage harvesting was obtained from *Stylosanthes scabra* 441. Mean plant height at forage harvesting was not significantly ($P>0.05$) different among the locations. The taller plant heights recorded for *Stylosanthes scabra* 441 genotypes resulted in better biomass yields. This is due to the fact that longer plants possess relatively more leaves and branches that may result in an increase in biomass yield [10].

Table 4: Mean plant height (cm) of four *Stylosanthes* genotypes tested across two locations/environments at forage harvesting stage.

| Genotypes | Location/Environments | | Combined analysis |
|----------------------------------|-----------------------|--------|-------------------|
| | Assosa | Kamash | |
| <i>Stylosanthes hamata</i> 167 | 40.39 | 37.64 | 39.02 |
| <i>Stylosanthes hamata</i> 75 | 44.66 | 38.13 | 41.39 |
| <i>Stylosanthes hamata</i> 15876 | 42.43 | 38.52 | 40.47 |
| <i>Stylosanthes scabra</i> 441 | 48.06 | 50.49 | 49.27 |
| Mean | 43.88 | 41.19 | 42.54 |
| CV | 18.68 | 20.97 | 19.17 |
| P-value | 0.4315 | 0.0528 | 0.0167 |

Forage dry matter yield

The forage dry matter yield of four tested *Stylosanthes* under two agro-ecological zones of Benishangul-Gumuz is indicated in Table 5. The result of each location and combined analysis showed that forage dry matter yield was non-significantly ($P>0.05$) different among the genotypes. The main effect difference among testing environments was significantly affect the forage dry matter yield ($P>0.05$). The highest forage dry matter yield was obtained at Assosa and this might be due to soil characteristics (Assosa soil is red, while Kamash soil is black). This might be due to the black soil being water logged which inhibits soil aeration, nutrient absorption and root growth that made plants stunted and reduced growth rate. The result also might be attributed to due to the leaf to stem ratio of the value

recorded for genotypes was higher at Kamash than Assosa testing environment. This leads to as the leaf to stem ratio increases, the forage dry matter yield decreases, however the nutrient content will be increase and this might be due the leaf part of the forage is more nutritious than stem. The overall mean (3.67 t ha⁻¹) for forge dry matter yield of the four *Stylosanthes* accessions in the present study was in line with the overall mean value (3.05 t ha⁻¹) reported by [16] for tested *Stylosanthes* accessions (for 166 *Stylosanthes hamata* and 34 *Stylosanthes scabra* accessions) tested on an acid soil, Soddo, southern Ethiopia [11,12].

Table 5: Mean forage DM yield (t/ha) of four *Stylosanthes* accessions tested across two locations/environments.

| Genotypes | Location/Environments | | Combined analysis |
|----------------------------------|-----------------------|--------|-------------------|
| | Assosa | Kamsh | |
| <i>Stylosanthes hamata</i> 167 | 4.1 | 1.99 | 3.04 |
| <i>Stylosanthes hamata</i> 75 | 5.1 | 2.05 | 3.58 |
| <i>Stylosanthes hamata</i> 15876 | 4.04 | 1.99 | 3.02 |
| <i>Stylosanthes scabra</i> 441 | 4.67 | 3.02 | 3.84 |
| Mean | 4.47 | 2.26 | 3.67 |
| CV | 30.04 | 28.68 | 60.99 |
| P-value | 0.8214 | 0.3203 | 0.7053 |

CONCLUSION

Stylosanthes accessions responded nonsignificant variations in forage dry matter yield across the testing environments, however, leaf to stem ratio showed significant variation among the accessions and *Stylosanthes scabra* 441 gave the highest leaf to stem ratio. Measured agronomic traits showed significant variations among the testing environments. The overall performance of *Stylosanthes* genotypes was better in Assosa than Kamash. This suggests that this location has better soil and climatic conditions for cowpea growing for forage purposes. Generally, *Stylosanthes scabra* 441 relatively showed the best performance for all measured agronomic traits across the testing environment and recommended for the study area and similar agro-ecologies.

ACKNOWLEDGEMENT

The field research was fully financed by Ethiopian Institute of Agricultural Research (EIAR). The authors would like to thank Assosa agricultural research center for facilitating the field research work.

AUTHORS' CONTRIBUTIONS

All authors contributed from the onset of the study and approved of the final version.

CONFLICT OF INTEREST DECLARATION

The authors have no conflict of interest to declare.

REFERENCES

1. Anele UY, Sudekum K, Aigbede OM, Welp G, Oni AO, Olanite JA, et al. Agronomic performance and nutritive quality of some

commercial and improved dual-purpose cowpea [*Vigna unguiculata* (L.) Walp.] varieties on marginal land in Southwest Nigeria. *Grassland Sci.* 2011;57:211-218.

- Sultan JI, Inam-Ur-Rahim, Nawaz H, Yaqoob M, Javed I. Nutritional evaluation of fodder trees leaves of northern grasslands of Pakistan. *Pak J Botany.* 2008;40:2503-2512.
- Hanson J, Heering JH. Genetic resources of *Stylosanthes* species. Proceeding of the Regional Workshop on the use of *Stylosanthes* in West Africa held in Kaduna, Nigeria, 1992, pp. 26-31.
- Chandra A. Diversity among *Stylosanthes* species: Habitat, edaphic and agro-climatic affinities leading to cultivar development. *J Environ Biol.* 2009;30(4):471-478.
- SAS. SAS user's guide: Statistics released 6.12. SAS Inc., Cary NC., USA, 2002.
- Basford KE, Cooper M. Genotype x environmental interactions and some considerations of their implications for wheat breeding in Australia. *Aust J Agric Res.* 1998;49:154-174.
- Kebede G, Feyissa F, Assefa G, Alemayehu M, Mengistu A, Kehalieu A, et al. Evaluation of Napier grass (*Pennisetum purpureum* (L.) Schumacher) accessions for agronomic traits under different environmental conditions of Ethiopia. *Int J Adv Res.* 2016;4(4): 1029-1035.
- Dixon AGO, Nukenine EN. Statistical analysis of cassava yield trials with the Additive Main Effects and Multiplicative Interaction (AMMI) model. *Afr J Root Tuber Crops.* 1997;3:46-50.
- Tudsri S, Jorgensen ST, Riddach P, Pookpakdi A. Effect of cutting height and dry season date on yield and quality of five Napier grass cultivars in Thailand. *Trop Grassland.* 2002;36:248-252.
- Smart AJ, Schacht WH, Moser LE, Volesky JD. Prediction of leaf/stem ratio using Near-Infrared Reflectance Spectroscopy (NIRS): A technical note. *Agron Horticult Faculty Publications.* 2004;39.
- Ceccarelli S. Adaptation to low/high input cultivation. Adaptation in plant breeding, Kluwer Academic Publishers, The Netherlands, 1997, pp. 225-236.
- Cameron D, Chakraborty S. Forage potential of *Stylosanthes* in different production systems. High-Yielding Anthracnose-Resistant *Stylosanthes* for Agricultural Systems. Australian Centre for International Agricultural Research (ACIAR), pp. 27-38.