

# Exploring Mycobacterial Food Infections: Unseen Culprits in the Culinary World

Hanshika Poorna\*

Department of Health Sciences, University of Hamburg, Hamburg, Germany

## INTRODUCTION

Foodborne illnesses are a global concern, affecting millions of individuals annually. While bacteria such as *Salmonella* and *E. coli* often take the spotlight, mycobacterial food infections represent a lesser-known but significant threat. This article explores mycobacterial food infection diseases, shedding light on the culprits, sources, clinical manifestations and preventive measures to enhance awareness and mitigate the risks associated with these infections.

## DESCRIPTION

### Mycobacteria in the culinary landscape

Mycobacteria are a diverse group of bacteria that includes the notorious *Mycobacterium tuberculosis*, responsible for tuberculosis and *Mycobacterium leprae*, the causative agent of leprosy. In addition to these well-known pathogens, several mycobacteria can cause foodborne infections. The most notable among them is *Mycobacterium avium* Subspecies Paratuberculosis (MAP), linked to Johne's disease in cattle and suspected to play a role in Crohn's disease in humans [1].

### Sources of contamination

Mycobacterial food infections are primarily associated with the consumption of unpasteurized dairy products, particularly milk and cheese. Infected animals, especially ruminants like cows, shed MAP in their feces, contaminating the environment. When raw milk or dairy products are consumed, there is a risk of ingesting viable mycobacteria [2].

In addition to dairy, mycobacteria can be present in various environmental sources, including water and soil. Contaminated water used in the production of fruits and vegetables or for washing food items can introduce mycobacteria into the food chain. Seafood, particularly shellfish, can also harbor mycobacterial species, posing a risk to consumers.

### Clinical manifestations

Mycobacterial food infections often present with non-specific symptoms, making diagnosis challenging. The incubation period can be prolonged, ranging from weeks to months, further complicating the identification of the source of infection. Common clinical manifestations include abdominal pain, diarrhea, weight loss and fatigue [3].

Johne's disease in cattle, caused by MAP, manifests as chronic enteritis, leading to weight loss, reduced milk production and ultimately death. In humans, mycobacterial food infections may present as a persistent gastrointestinal illness resembling inflammatory bowel diseases, such as Crohn's disease.

### Diagnostic challenges

The diagnosis of mycobacterial food infections is intricate due to the slow growth of these bacteria and the similarity of symptoms to other gastrointestinal disorders. Laboratory tests, including culture and molecular methods, are employed to detect and identify mycobacterial species. However, the slow growth characteristics of these bacteria can delay diagnosis, impacting the timely initiation of appropriate treatment.

### Preventive measures

Preventing mycobacterial food infections involves a combination of regulatory measures, hygiene practices and consumer awareness. Pasteurization of dairy products is a critical step in eliminating mycobacterial contamination. Strict adherence to hygiene protocols in the handling of food, especially in food production and processing facilities, is essential to prevent cross-contamination [4].

Regulatory bodies play a pivotal role in enforcing standards for the production and distribution of food products. Regular monitoring of dairy farms, water sources and food processing facilities can help identify and mitigate potential sources of mycobacterial contamination. Public health campaigns aimed at educating consumers about the risks associated with consuming

**Correspondence to:** Hanshika Poorna, Department of Health Sciences, University of Hamburg, Hamburg, Germany; E-mail: Aakhus.hans@mail.in

**Received:** 05-Dec-2023, Manuscript No. MDTL-23-28357; **Editor assigned:** 07-Dec-2023, PreQC No. MDTL-23-28357 (PQ); **Reviewed:** 21-Dec-2023, QC No. MDTL-23-28357; **Revised:** 13-Jan-2025, Manuscript No. MDTL-23-28357 (R); **Published:** 20-Jan-2025, DOI: 10.35248/2161-1068.25.15.537

**Citation:** Poorna H (2025) Exploring Mycobacterial Food Infections: Unseen Culprits in the Culinary World. *Mycobact Dis*. 15:537.

**Copyright:** © 2025 Poorna H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

raw or unpasteurized dairy products can contribute to preventive efforts.

### Global implications and emerging research

Mycobacterial food infections have global implications, affecting both developed and developing nations. The interconnectedness of the food supply chain and international trade heightens the risk of the spread of these infections across borders. Surveillance systems and collaboration between countries are crucial for monitoring and responding to mycobacterial foodborne outbreaks.

Emerging research in this field focuses on understanding the prevalence of mycobacteria in various food sources, elucidating their survival mechanisms and developing rapid diagnostic tools. Advances in genomics and molecular techniques enable researchers to explore the genetic diversity of mycobacterial strains, shedding light on their adaptation to different environments and host species.

### Future challenges and opportunities

As our understanding of mycobacterial food infections deepens, challenges and opportunities emerge on the horizon. Improved diagnostic technologies, increased surveillance and enhanced communication between researchers, healthcare professionals and regulatory bodies are essential components of an effective strategy against these infections [5].

Public health interventions should not only target the prevention of mycobacterial foodborne illnesses but also address the challenges associated with their diagnosis and treatment. Multidisciplinary collaboration, involving microbiologists,

epidemiologists, clinicians and policymakers, is crucial for developing comprehensive strategies to tackle these infections on a global scale.

## CONCLUSION

Mycobacterial food infections, though less recognized than some other foodborne illnesses, pose a significant threat to public health. The complexities associated with their diagnosis and the non-specific nature of symptoms make these infections challenging to detect and manage. By fostering awareness, implementing preventive measures and investing in research, we can enhance our ability to mitigate the risks posed by mycobacterial foodborne pathogens and safeguard the well-being of consumers worldwide.

## REFERENCES

1. Ayoola MB, Pillai N, Nanduri B, Rothrock MJ, Ramkumar M. Preharvest environmental and management drivers of multidrug resistance in major bacterial zoonotic pathogens in pastured poultry flocks. *Microorganisms*. 2022;10(9):1703.
2. Jovanovic J, Ornelis VF, Madder A, Rajkovic A. *Bacillus cereus* food intoxication and toxicoinfection. *Compr Rev Food Sci Food Saf*. 2021;20(4):3719-3761.
3. Paxson H, Helmreich S. The perils and promises of microbial abundance: Novel natures and model ecosystems, from artisanal cheese to alien seas. *Soc Stud Sci*. 2014;44(2):165-193.
4. Wilson ME. The traveller and emerging infections: Sentinel, courier, transmitter. *J Appl Microbiol*. 2003;94(S1):1-11.
5. Brody AL, Bugusu B, Han JH, Sand CK, McHugh TH. Innovative food packaging solutions. *J Food Sci*. 2008;73(8):107-116.