

## Survey on Bandage Contact Lens Practice in the United Kingdom

Philipp Baback Baenninger<sup>1,2\*</sup>, Christiana Dinah<sup>1</sup> and Francisco Carlos Figueiredo<sup>1</sup>

<sup>1</sup>Department of Ophthalmology, Royal Victoria Infirmary, Newcastle University, Newcastle upon Tyne, UK

<sup>2</sup>Department of Ophthalmology, Cantonal Hospital of Lucerne, Lucerne, Switzerland

### Abstract

**Purpose:** To determine the opinion regarding the prescribing practices of bandage contact lenses (BCL) amongst members of the Bowman Club (UK Cornea Society).

**Methods:** In June 2011, followed by a reminder in July, a questionnaire was sent out to all 128 members of the Bowman Club. The survey included 19 questions regarding indications, preferred type, methods of insertion, concomitant medication, complications related to BCL use among other questions.

**Results:** The survey was completed anonymously by 52 (40.6%) consultant ophthalmologists either online (88.5%) or by post (11.5%). The most common indication is pain relief by 51 (98%), followed by promotion of epithelial healing by 49 (94.2%) respondents. Silicon hydrogel soft contact lens is the most commonly used BCL by 39 (75%) consultants.

There was a higher incidence of secondary corneal ulcers reported if consultants were using non-sterile (51.9%) versus sterile (15.4%) insertion technique as well as use (26.9%) versus no use of topical prophylactic antibiotic (40.4%).

**Conclusions:** This is the first survey on practice pattern of BCL use amongst consultant ophthalmologists with expertise in the management of a subspecialty in Ocular Surface Diseases in the UK. It demonstrates that the most common indication for BCL use is pain relief, with silicone hydrogel soft contact lenses being the most frequently used. Secondary corneal ulcers were more frequently seen by consultants that neither use sterile insertion technique nor prophylactic topical antibiotics.

**Keywords:** Survey; Bandage contact lenses; Practice patterns; Secondary corneal ulcer

### Introduction

Although bandage contact lenses (BCL) have been in widespread use in the United Kingdom (UK) [1] for many years, limited data is available on the prescribing practice amongst UK Consultants with subspecialty interest in Ocular Surface Diseases, members of the Bowman Club (UK Cornea Society).

Therapeutic BCL helps to maintain ocular tissue integrity [1] and are used for pain relief [2-7], mechanical protection of the ocular surface [2-4,8,9] and promotion of epithelial healing [10,11]. A barrier effect of therapeutic contact lenses to airborne antigen was identified in ocular allergy [12].

There have been multiple reports about safety and efficacy of BCL for different indications as in recurrent erosion syndrome [13-15], bullous keratopathy [16,17] and immediately after LASEK [18] or phototherapeutic keratectomy [19] in adults and in children [20]. Large-diameter BCL has also been reported as a useful treatment option in the management of refractory vernal ulcers [21]. Gas-permeable scleral contact lenses provide an additional effective management option in severe ocular surface disease (e.g. Stevens-Johnson syndrome) [22]. As the later two uses of BCL are limited to specific indications and high severity of ocular surface disease, they were not assessed in this survey.

Moutray et al. [14] found that BCL were normally well tolerated by patients. Silicone hydrogel materials provide clinicians with a potential alternative to hydrogel BCL [23] as Silicon hydrogel BCL provide additional mechanical protection, safer overnight wear, increased oxygen transmission [24] and relatively low water content [14,25].

Although BCL wear is considered to be safe, complications secondary to BCL use are increasingly noted in patients who are wearing it for longer period [26-31].

The role of prophylactic topical antibiotics to prevent complications is unclear [32]. Ozkan et al. [33] recently showed that topical antibiotics were well tolerated, but there was no significant difference in the numbers and types of microbes recovered from lens sample either treated prophylactically with tobramycin 0.3% or placebo eye drops.

No reports on best insertion technique or concomitant use of pharmaceuticals can be found.

A survey in North America [23] revealed that BCL were most commonly used for corneal wound healing and managing post-operative complications, however this report had a very poor response rate of 3.4%, limiting the reliability of their results. Therefore, the aim of the study was to determine the opinion regarding the prescribing practices of BCL in the UK amongst members of the Bowman Club.

### Methods

In June 2011, followed by a reminder in July 2011, a questionnaire (Supplementary file) was sent by email to all 128 members of the Bowman Club. In order to become part of the Bowman Club, members need to hold a consultancy position in the UK and have a subspecialty interest in Ocular Surface Diseases. Name and email addresses of all members are stored by the Bowman Club administration.

**\*Corresponding author:** Dr. Philipp B Baenninger, Augenklinik, Cantonal Hospital of Lucerne, CH-6000 Lucerne-16, Switzerland, Tel: 41 76 438 5153; Fax: +41 205 3421; E-mail: [philipp.baenninger@luks.ch](mailto:philipp.baenninger@luks.ch)

**Received** December 03, 2013; **Accepted** February 10, 2014; **Published** February 19, 2014

**Citation:** Baenninger PB, Dinah C, Figueiredo FC (2014) Survey on Bandage Contact Lens Practice in the United Kingdom. J Clin Exp Ophthalmol 5: 325. doi: 10.4172/2155-9570.1000325

**Copyright:** © 2014 Baenninger PB, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The invitation email to all members included a link to participate in the survey online as well as an attached PDF copy of the questionnaire that could be printed, completed and returned by mail. The survey included 19 questions regarding indications, preferred type, methods of insertion, concomitant medication, complications related to BCL use among other questions. The survey was then returned anonymously by post or online to the study group. Participants in the survey did not receive monetary reimbursement for their participation.

Statistical analysis was performed in order to determine the significance of the association between complications and methods of insertion, using Fisher's exact test.

## Results

The survey was completed anonymously by 52 (40.6%) consultant ophthalmologists either online (88.5%) or by post (11.5%). The majority of consultant ophthalmologist with a subspecialty interest in ocular surface diseases had been working for more than 10 years (n=31, 59.6%), whereas 14 (26.9%) were practicing for 5-10 years and 7 (13.5%) for less than 5 years as consultant.

## Indication

The most common indication was pain relief (e.g. bullous keratopathy or recurrent erosion syndrome) by 51 (98%), followed by promotion of epithelial healing (e.g. persistent epithelial defects, recurrent erosion syndrome), by 49 (94.2%) respondents. Furthermore, 43 (82.7%) of members use BCL for postoperative protection (e.g. following conjunctival/corneal surgery); 40 (76.9%) for apposition of wound edges (e.g. corneal perforation, after suture removal); 38 (73.1%) for mechanical protection of ocular surface (e.g. trichiasis, entropion) and 24 (46.2%) for maintenance of ocular hydration (e.g. dry eye, corneal exposure) (Table 1).

## Type of BCL

Silicon hydrogel soft contact lens is the most commonly used BCL by 39 (75%) consultants among them, PureVision® being the most frequently used by 19 consultants (37.3%) followed by ACUVUE® by 6 (11.8%) and NIGHT&DAY® by 5 (9.8%) consultants. High-water content hydrogel soft contact lens is used by 29 (55.8%) consultants, among them Proclear being the most frequently used by 3 consultants (5.8%) followed by Precision UV by 2 (3.8%) and Permalens by 1 (1.9%) consultant. Nine (17.3%) consultants reported no preference for a particular BCL. No trend was detected for the use of a specific BCL type for a specific indication.

## Handling of BCL

39 (75%) consultants insert the BCL manually (e.g. fingertips)

Indication	Consultant n(%)
Pain relief (e.g. bullous keratopathy, recurrent erosion syndrome)	51 (98%)
Promotion of epithelial healing (e.g. persistent epithelial defects, recurrent erosion syndrome)	49 (94.2%)
Postoperative protection (e.g. following conjunctival/corneal surgery)	43 (82.7%)
Apposition of wound edges (e.g. corneal perforation, after suture removal)	40 (76.9%)
Mechanical protection of ocular surface (e.g. trichiasis, entropion)	38 (73.1%)
Maintenance of ocular hydration (e.g. dry eye, corneal exposure)	24 (46.2%)

Table 1: Indications for Bandage contact lens use.

and 19 (25%) use sterile forceps technique (or equivalent). BCL replacement protocol is at monthly (48.1%) or six-weekly (19.2%) time interval, but varies according to 25 (48.1%) respondents on the underlying diagnosis (range: fortnightly to quarterly). Replacement takes place in the consultant Ocular Surface Disease clinic (70.6%) and/or optometry department (45.1%), rarely in the community (3.9%). Insertion by optometry department protocol is manually (71.2%) and with sterile forceps or equivalent, e.g. using the nozzle of Minims by suction (13.5%) and the usual patient review by medical staff in this setting is performed three-monthly (37.5%), more than four-monthly (18.8%) or varies (31.3%). Punctum plugs are not routinely inserted (98.0%) in combination with BCL.

## BCL policy in paediatric patients

22 consultants (42.3%) change their BCL in paediatric patients compared to 18 (34.6%) consultants, who follow the same policy as in adults. Twelve (23.1%) consultants do not treat paediatric patients in their clinic. The BCL policy changes as followed: 17 (32.7%) consultants never prescribe BCL in children, 3 (5.8%) perform BCL replacement more often, 1 (1.9%) replaces BCL only in consultant's clinic and not by the optometry department and 1 (1.9%) uses BCL only for specific indications, e.g., corneal protection.

## Concomitant use of pharmaceuticals

All consultants are happy to use concomitant topical treatment for the underlying condition; including antibiotics by 52 (100%), artificial tears by 47 (90.4%), steroids by 45 (86.5%), anti-glaucoma by 32 (61.5%), cycloplegics by 30 (57.7%), non-steroidal anti-inflammatory by 12 (23.1%) and anti-virals by 8 (15.4%). Twenty two consultants responded (42.3%) usually prescribe prophylactic topical antibiotics such as G. Chloramphenicol (95.5%). However, 18 (34.6%) only use preservative-free medication, 25 (48.1%) use preservative-free medication if available and 9 (17.3%) do not use preservative-free eye drops.

## Complications

Infected corneal ulcer secondary to BCL use is reported by 67.3% consultants: 1-3 ulcers/year by 48.1%, 4-6 by 15.4%, and >10 by 3.8% in the last 12 months. Majority of corneal ulcers are of bacterial (97.6%) aetiology, one consultant (2.4%) reported a fungal corneal ulcer secondary to BCL use. There was a higher incidence of secondary corneal ulcers reported if consultants were using non-sterile (51.9%) versus sterile (15.4%) insertion technique, although it was not significant (p=0.743) as well as non use of topical prophylactic antibiotic (40.4%) versus the use (26.9%) of topical prophylactic antibiotic (p=0.776).

## Discussion

This is the first survey on practice pattern of BCL use amongst consultant ophthalmologists in the UK with a subspecialty interest in ocular surface diseases. It demonstrates that the most common indication for BCL use is pain relief, with silicone hydrogel soft contact lenses being the most frequently used. A high incidence of secondary corneal ulcers per year is reported, and it is important to noticed that topical prophylactic antibiotic management is only used by 42.3% consultants and 75% report non-sterile BCL insertion technique. Secondary corneal ulcers were more frequently seen with non-sterile than sterile insertion technique, use of prophylactic topical antibiotics resulted in less secondary corneal ulcers.

Amongst many possible indications for BCL, in this survey the most commonly reported indications are pain relief, promotion of epithelial healing and postoperative protection. Similar results were reported by Karlsgard [23] and Montero [34]. In contrast, silicone hydrogel BCL PureVision® is most popular amongst consultants in the UK, followed by ACUVUE® and NIGHT&DAY®, which are most frequently used in the US and Canada [23].

Unfortunately, no reports on insertion technique or concomitant use of pharmaceuticals can be found in the literature, therefore, this is the first study reporting two possible risk factors of BCL usage in the UK, i.e., non-sterile insertion technique and no use of prophylactic topical antibiotics that can lead to a higher incidence of secondary corneal ulcers.

Using sterile insertion technique is expected to reduce the risk of contamination, especially as BCL are indicated in patients with compromised ocular surface that are more prone for infection. Although, the role of prophylactic topical antibiotics is unclear, Ozkan [33] reported that instillation of tobramycin 0.3% twice daily did not reduce contamination rate of worn lenses and did not result in significant changes in numbers and profiles of microorganisms colonized on the lens surface over a 3-month study period when compared to a placebo group. The study also showed a rate of 69% of sterile lenses in continuous lens wear. In comparison, Keay [35] reported only 28% sterile lenses when no eye drops were used, this results suggested that prophylactic antibiotics might reduce the risk of BCL contamination. Kalayci [36] demonstrated that ciprofloxacin-pre-soaked contact lenses retained significant antimicrobial activity up to 12 hours and Fraunfelder [13] regards the use of ofloxacin twice daily as a prophylactic dose when compared to four times daily. Therefore, based on our results further investigation regarding the role of prophylactic topical antibiotics is needed.

A strength of this survey is the good response rate of 40.6% amongst members of the Bowman Club, who are consultant ophthalmologist with a subspecialty interest in ocular surface diseases. Furthermore, most of respondents have considerable expertise working in the field for more than 10 years, therefore a good representation of the ophthalmological community at large.

It is inherent to this survey, that the results of complications secondary to BCL use should be regarded as a trend, however, our results may suggest a review of BCL practice by all ophthalmologists using BCL as part of their routine care to include sterile lens insertion using forceps and the use of prophylactic topical antibiotic concomitantly with BCL wear.

General recommendations drawn from this survey may include the use of BCL for indications like pain relief, promotion of epithelial healing and postoperative protection. In order to reduce the risk of secondary corneal ulcer, regular, monthly replacement of BCL, the use of sterile insertion technique and prophylactic topical antibiotics should be considered.

#### Acknowledgment

The authors thank and acknowledge all Members of the Bowman Club (UK Cornea Society) for their invaluable contribution.

#### References

1. Ehrlich DP (2006) Therapeutic contact lenses. [Continuing Education and Training]. July 14. Accessed June 21, 2013.
2. McDermott ML, Chandler JW (1989) Therapeutic uses of contact lenses. *Surv Ophthalmol* 33: 381-394.

3. Gasset AR, Kaufman HE (1970) Therapeutic uses of hydrophilic contact lenses. *Am J Ophthalmol* 69: 252-259.
4. Smiddy WE, Hamburg TR, Kracher GP, Gottsch JD, Stark WJ (1990) Therapeutic contact lenses. *Ophthalmology* 97: 291-295.
5. Leibowitz HM (1972) Hydrophilic contact lenses in corneal disease. IV. Penetrating corneal wounds. *Arch Ophthalmol* 88: 602-606.
6. Mondino BJ, Zaidman GW, Salamon SW (1982) Use of pressure patching and soft contact lenses in superior limbic keratoconjunctivitis. *Arch Ophthalmol* 100: 1932-1934.
7. Forstot SL, Binder PS (1979) Treatment of Thygeson's superficial punctate keratopathy with soft contact lenses. *Am J Ophthalmol* 88: 186-189.
8. Andrew NC, Woodward EG (1989) The bandage lens in bullous keratopathy. *Ophthalmic Physiol Opt* 9: 66-68.
9. Aquavella JV (1974) Editorial: Therapeutic uses of hydrophilic lenses. *Invest Ophthalmol* 13: 484-486.
10. Hykin PG, Foss AE, Pavesio C, Dart JK (1994) The natural history and management of recurrent corneal erosion: a prospective randomised trial. *Eye (Lond)* 8: 35-40.
11. Williams R, Buckley RJ (1985) Pathogenesis and treatment of recurrent erosion. *Br J Ophthalmol* 69: 435-437.
12. Wolffsohn JS, Emberlin JC (2011) Role of contact lenses in relieving ocular allergy. *Cont Lens Anterior Eye* 34: 169-172.
13. Fraunfelder FW, Cabezas M (2011) Treatment of recurrent corneal erosion by extended-wear bandage contact lens. *Cornea* 30: 164-166.
14. Moutray TN, Frazer DG, Jackson AJ (2011) Recurrent erosion syndrome—the patient's perspective. *Cont Lens Anterior Eye* 34: 139-143.
15. Langston RH, Machamer CJ, Norman CW (1978) Soft lens therapy for recurrent erosion syndrome. *Ann Ophthalmol* 10: 875-878.
16. Ruben M (1975) Soft contact lens treatment of bullous keratopathy. *Trans Ophthalmol Soc U K* 95: 75-78.
17. Lim N, Vogt U (2006) Comparison of conventional and silicone hydrogel contact lenses for bullous keratoplasty. *Eye Contact Lens* 32: 250-253.
18. Orucov F, Frucht-Pery J, Raikup FD, Strasman E, Landau D, et al. (2010) Quantitative assessment of bandage soft contact lens wear immediately after LASIK. *J Refract Surg* 26: 744-748.
19. Seiler T, McDonnell PJ (1995) Excimer laser photorefractive keratectomy. *Surv Ophthalmol* 40: 89-118.
20. Bendoriene J, Vogt U (2006) Therapeutic use of silicone hydrogel contact lenses in children. *Eye Contact Lens* 32: 104-108.
21. Quah SA, Hemmerdinger C, Nicholson S, Kaye SB (2006) Treatment of refractory vernal ulcers with large-diameter bandage contact lenses. *Eye Contact Lens* 32: 245-247.
22. Romero-Rangel T, Stavrou P, Cotter J, Rosenthal P, Baltatzis S, et al. (2000) Gas-permeable scleral contact lens therapy in ocular surface disease. *Am J Ophthalmol* 130: 25-32.
23. Karlgard CC, Jones LW, Moresoli C (2004) Survey of bandage lens use in North America, October-December 2002. *Eye Contact Lens* 30: 25-30.
24. Alvord L, Court J, Davis T, Morgan CF, Schindhelm K, et al. (1998) Oxygen permeability of a new type of high Dk soft contact lens material. *Optom Vis Sci* 75: 30-36.
25. Rubinstein MP (2003) Applications of contact lens devices in the management of corneal disease. *Eye (Lond)* 17: 872-876.
26. Stapleton F, Keay L, Edwards K, Naduvilath T, Dart JK, et al. (2008) The incidence of contact lens-related microbial keratitis in Australia. *Ophthalmology* 115: 1655-1662.
27. Kaufman HE (1979) Problems associated with prolonged wear soft contact lenses. *Ophthalmology* 86: 411-417.
28. Dart JK, Radford CF, Minassian D, Verma S, Stapleton F (2008) Risk factors for microbial keratitis with contemporary contact lenses: a case-control study. *Ophthalmology* 115: 1647-1654, 1654.
29. Stapleton F, Keay L, Jalbert I, Cole N (2007) The epidemiology of contact lens related infiltrates. *Optom Vis Sci* 84: 257-272.

- 
30. Ormerod LD, Smith RE (1986) Contact lens-associated microbial keratitis. *Arch Ophthalmol* 104: 79-83.
  31. Keay L, Stapleton F, Schein O (2007) Epidemiology of contact lens-related inflammation and microbial keratitis: a 20-year perspective. *Eye Contact Lens* 33: 346-353, discussion 362-3.
  32. Murali K, Aasuri S (1997) Bandage Contact Lenses in Ocular Disorders. *International Contact Lense Clinic* 24: 207-212.
  33. Ozkan J, Zhu H, Gabriel M, Holden BA, Willcox MD (2012) Effect of prophylactic antibiotic drops on ocular microbiota and physiology during silicone hydrogel lens wear. *Optom Vis Sci* 89: 326-335.
  34. Montero J, Sparholt J, Mely R (2003) Retrospective case series of therapeutic applications of a lotrafilcon A silicone hydrogel soft contact lens. *Eye Contact lens* 29(1 Suppl): S54-S56.
  35. Keay L, Willcox MD, Sweeney DF, Morris CA, Harmis N, et al. (2001) Bacterial populations on 30-night extended wear silicone hydrogel lenses. *CLAO J* 27: 30-34.
  36. Kalayci D, Basci N, Kortunay S, Hasiripi H, Bozkurt A (1999) Penetration of topical ciprofloxacin by presoaked medicated soft contact lenses. *CLAO J* 25: 182-184.