

Bionomics of Rose Thrips, *Frankliniella Occidentalis* Pergande

Kanara HG* and Acharya MF

Department of Entomology, College of Agriculture, Junagadh Agricultural University, Junagadh, Gujarat, India

*Corresponding author: Kanara HG, Department of Entomology, College of Agriculture, Junagadh Agricultural University, Junagadh, Gujarat, India, Tel: 9558891750; E-mail: kanarahiren1@gmail.com

Rec date: Apr 15, 2014; Acc date: Aug 26, 2014; Pub date: Aug 30, 2014

Copyright: © 2014 Kanara HG, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

A lab trial was conducted during 2012-13 at Department of Entomology, College of Agriculture, JAU, Junagadh to study the bionomics of rose thrips, *Frankliniella occidentalis* Pergande. Laboratory studies revealed that the female laid their eggs in the leaf tissues by sharp ovipositor. The average incubation period was found 5.28 day. The larva passed through two distinct instar and average duration of each instar was 1.72 and 4.76 day, respectively. The total larval period was on an average of 6.48 day with the total pupal period of 3.96 day. The average pre-oviposition, oviposition and post-oviposition period were 3.12, 32.68 and 4.20 day, respectively. The number of progenies produced by a single female varied from 28 to 69 with an average of 58.8. The average longevity of male and female was 21.60 and 42.36 day with entire life span of 25.52 and 48.40 day, respectively. The sex ratio of male to female was worked out as 1: 1.73 at a constant temperature of $25 \pm 1^\circ\text{C}$.

Keywords: Bionomics; Rose; *Frankliniella occidentalis*

Introduction

Western flower thrips, *Frankliniella occidentalis* (Pergande) [Thripidae: Thysanoptera] is a serious problem on rose grown under protected cultivation. Rose (*Rosa* sp.) is one of the nature's beautiful creations and is universally called as 'queen of flower'. Cultivation of rose under protected conditions has gained importance in recent years due to its export potential. Rose is used for worship, in making garlands and bouquets [1]. The pest feeding results in mottling, severe curling, browning and drying of tender leaves, sepals, tender stalks, outer petals of green and half opened buds, which turn to brown colour and appear as if they are burnt. Damaged flowers get discoloured and distorted in shape and reduced in size [2]. This pest can cause 28-95 percent damage with a population density of 11-33 thrips/flower [3]. At present the information on bionomics of this pest is scarce. Hence, the study was conducted under laboratory conditions and the findings of the present study would help in managing this pest.

Materials and Methods

The present investigation on bionomics of thrips, *F. occidentalis* on rose was carried out at Department of Entomology, College of Agriculture, Junagadh Agricultural University, Junagadh during 2012-13. The study was carried out at a constant temperature of $25 \pm 1^\circ\text{C}$ by using B.O.D. incubator. In order to develop the initial culture of thrips, *F. occidentalis*, large number of adults were collected with the help of aspirator from the rose plants, five females and two males were picked up individually by means of moistened camel hair brush and released gently into a glass vial (3 cm × 1 cm) held in an inverted position. A young leaflet of rose was then introduced into glass vial and the vial was closed with bark cork. Thus, field collected adults were distributed in 25 vials to obtain large number of progenies. The vials were kept in an incubator adjusted to $25 \pm 1^\circ\text{C}$ temperature for oviposition. As soon as the larvae emerged out from the leaflets, they

were reared separately into the glass vials. The leaflets of rose were changed once every 2 days until the larvae pupated.

The rearing was continued till the emergence of adults and they were used for further investigations. The sexes of adults were determined on the basis of their body colour, size and abdominal tip. The males were smaller in size, pale in colour with rounded abdominal tip, whereas the females were dark brown to black with pointed abdominal tip. The adults, thus, obtained were paired for the further study. Each stage of the thrips were observed and measured in Leica microscope at Biocontrol laboratory of Entomology.

Egg

A pair of male and female was enclosed in a glass vial (3 cm×1 cm) containing a rose leaflet for oviposition. The leaflet was collected after 24 hours from the vial and kept individually in a separate vial for hatching of eggs. The procedure was continued till the death of female. The incubation period and number of larvae emerged out was recorded.

Larva

With a view to study the larval instars and their duration, the newly emerged young larva was transferred with a help of a wet camel hair brush and kept individually in a glass vial containing rose leaflet, which was changed daily in the morning. The individual leaflet was examined critically under binocular microscope to confirm the moulting. The observations on number of instars, instar duration and total larval duration were recorded.

Pupa

For ease of pupation, a thin layer of fine soil was kept at bottom of the glass vial as the pupation take place in the soil. The pupa when formed was collected and kept individually into glass vial for emergence of adults. The prepupal and pupal periods were recorded.

Adult

The adults emerged from the pupae were paired on the same day. Each pair was introduced into glass vial to study the fecundity and longevity. Fresh rose leaflets were provided into glass vials for food and oviposition. The leaflet was changed at an interval of 24 hours. The old leaflets were collected and kept into another glass vials for emergence of progenies. Observations on pre-oviposition, oviposition and post-oviposition and longevity of adults were recorded.

Results and Discussion

Present studies revealed that the female of *F. occidentalis* inserted their eggs, partially or completely, into parenchyma tissue of leaves or flower by making incision with the help of saw like ovipositor. Since, the eggs were embedded into leaf tissues, they could not be studied.

However, the number of eggs laid by a female, incubation period, pre and post ovipositional periods were derived on the basis of number of larvae emerged out from the eggs.

Incubation period of *F. occidentalis* varied from 4 to 7 day with an average of 5.28 ± 0.73 day. This finding partially supports the results of Ronald and Jayma [4], who observed the incubation period as 6.37 day on rose at a constant temperature of 20°C. The larva passed through two distinct instar before attaining the pupal stage. The newly emerged first instar larva was white or nearly transparent which later on turned to yellow, orange, crimson or purple in colour. Its body consisted of a head; 3 thoracic segments and 11 abdominal segments with 3 pair of similarly structured legs and no wing buds. When the first instars larvae become doubled in size, they searched for a niche to molt. Similar observations were also reported by Lewis [5], when the pest was reared on rose in London.

| Sr. No. | Different periods (Days) | No. of observations | Minimum | Maximum | Av. \pm S.D. |
|---------|----------------------------|---------------------|---------|---------|------------------|
| 1 | Incubation period | 25 | 4 | 7 | 5.28 ± 0.73 |
| 2 | First larval instar | 25 | 1 | 2 | 1.72 ± 0.45 |
| 3 | Second larval instar | 25 | 3 | 6 | 4.76 ± 0.72 |
| 4 | Total larval period | 25 | 4 | 8 | 6.48 ± 1.17 |
| 5 | Pre pupa period | 25 | 1 | 3 | 1.76 ± 0.52 |
| 6 | Pupa period | 25 | 2 | 3 | 2.20 ± 0.57 |
| 7 | Total pupal period | 25 | 3 | 6 | 3.96 ± 1.09 |
| 8 | Pre-oviposition period | 25 | 2 | 4 | 3.12 ± 0.49 |
| 9 | Oviposition period | 25 | 28 | 34 | 32.68 ± 1.18 |
| 10 | Post-oviposition period | 25 | 3 | 5 | 4.20 ± 0.64 |
| 11 | Longevity of male | 25 | 19 | 23 | 21.60 ± 0.86 |
| 12 | Longevity of female | 25 | 41 | 44 | 42.36 ± 0.75 |
| 13 | Entire life span of male | 25 | 19 | 31 | 25.52 ± 1.98 |
| 14 | Entire life span of female | 25 | 39 | 52 | 48.40 ± 2.18 |

Table 1: Duration of different developmental stages of *F. occidentalis* on rose

In the present study, the body length of first and second larval instar (Table 2) varied from 0.54 to 0.57 mm and 0.85 to 0.88 mm with an average of 0.56 ± 0.01 mm and 0.86 ± 0.01 mm, respectively. The duration of first and second larval instars varied from 1 to 2 day and 3 to 6 day with an average of 1.72 ± 0.45 and 4.76 ± 0.72 day, respectively. The total larval period varied from 4 to 8 day with an average of 6.48 ± 1.17 day. Almost similar result was obtained by

Ronald and Jayma [6] on rose. Two pupal stages were recorded in the present study. Both stages had functional legs. Before pupation, the larva entered into prepupal stage which was somewhat darker than the second instar larva. Prepupal stage had wing buds, rudimentary antennae and did not excrete. The pupa had developed antennae that curve back over the head, elongated wing pads and legs. Similar observations were recorded by Lewis [5].

| Sr. No. | Different stages | No. of observation | Length (mm) | | Av. \pm S.D. |
|---------|----------------------|--------------------|-------------|---------|-----------------|
| | | | Minimum | Maximum | |
| 1 | First larval instar | 25 | 0.54 | 0.57 | 0.56 ± 0.01 |
| 2 | Second larval instar | 25 | 0.85 | 0.88 | 0.86 ± 0.01 |
| 3 | Pre pupa | 25 | 0.76 | 0.78 | 0.77 ± 0.02 |

| | | | | | |
|---|--------|----------------------------------|------|------|-------------|
| 4 | Pupa | 25 | 0.79 | 0.82 | 0.81 ± 0.01 |
| 5 | Male | 25 | 0.96 | 1.11 | 1.01 ± 0.08 |
| 6 | Female | 25 | 1.20 | 1.50 | 1.40 ± 0.10 |
| | | Breadth with wing expansion (mm) | | | |
| 7 | Male | 25 | 1.49 | 1.54 | 1.53 ± 0.03 |
| 8 | Female | 25 | 1.89 | 1.94 | 1.92 ± 0.04 |

Table 2: Measurements of different stages of *F. occidentalis*

Studies on body length of pre pupa revealed that it ranged from 0.76 to 0.78 mm with an average of 0.77 ± 0.02 mm, while the length of pupa ranged from 0.79 to 0.82 mm with an average of 0.81 ± 0.01 mm. They were presented in Table 2. Present studies on pupal period indicated that the pre pupal and pupa period varied from 1 to 3 day and 2 to 3 day with an average of 1.76 ± 0.52 and 2.20 ± 0.57 day, respectively. The total period varied from 3 to 6 day with an average of 3.96 ± 1.09 day. The duration of different developmental stages of *F. occidentalis* were presented in Table 1. Total 25 number of individual were observed for the study of each duration of developmental stages.

Adult was thin, elongated and dorso ventrally flattened with light yellow in colour. Female was larger than the male with light brown patches on abdomen. Adult possessed two pair of long and slender membraneous wings usually with only a few longitudinal setae bearing veins which were covered with minute microtrichia. The forewings covered hindwings and abdomen. It is evident from the result that body length of the male thrips ranged from 0.96 to 1.11 mm (Average 1.01 ± 0.08 mm) in length and 1.49 to 1.54 mm (Average 1.53 ± 0.03 mm) in breadth, whereas the female thrips measured from 1.20 to 1.50 mm with an average of 1.40 ± 0.10 mm in length and 1.89 to 1.94 mm

with an average of 1.92 ± 0.04 mm in breadth. Kirk and Terry (6) reported that the adult male was about 1 mm long and female was slightly larger, about 1.4 mm in length. Present study indicated that male and female thrips lived for 19 to 23 day (Average 21.60 ± 0.86 day) and from 41 to 44 day (Average 42.36 ± 0.75 day), respectively.

The results revealed the pre-oviposition, oviposition and post-oviposition periods varied from 2 to 4, 28 to 34 and 3 to 5 day with an average of 3.12 ± 0.49 , 32.68 ± 1.18 and 4.20 ± 0.64 day, respectively. A single female produced 28 to 69 progenies with an average of 58.8 ± 6.78 . Kirk and Terry (6) observed that each female may lay 40 to over 100 eggs in the tissue of the plant, often in the flower.

The sex ratio of male and female of *F. occidentalis* was 1:1.73 presented in Table 3. The entire life span of male varied from 19 to 31 day with an average of 25.52 ± 1.98 day and that of female, 39 to 52 day with an average of 48.40 ± 2.18 day. The slightly variations observed in the present investigation with earlier findings in the bionomics of *F. occidentalis* may be due to the changes in the environmental factors.

| No. of adults observed | Sex | | Sex-ratio (Male: Female) |
|------------------------|-------------|---------------|-----------------------------|
| | No. of male | No. of female | |
| 200 | 73 | 127 | 1 : 1.73 |

Table 3: Sex ratio of *F. occidentalis*

References

- Bose TK, Yadav, LP (1989) Commercial flowers. Naya Prakash, Calcutta.
- Jhansi Rani, B, Jagan Mohan N (1997) Pest management in Horticultural crops. In: Progressive Floricultural. Yadav IS and Chaudhary ML (Eds), House of Span, pp: 169-181
- Gahukar RT (2003) Factors influencing thrips abundance and distribution on rose flowers in central India. Journal of the Entomological Research Society 27: 271-279.
- Ronald FL, Jayma L (1993) Annual research report, Department of Entomology, Honolulu, Hawaii 11-14.
- Lewis T (1973) Thrips their biology, ecology and economic importance. Academic Press: London, New York.
- Kirk DJ, Terry IL (2003) The spread of the western flower thrips *Frankliniella occidentalis* (Pergande). Agricultural and Forest Entomology 5: 301-310.