

## Hydroxyurea Therapy in Patients with Sickle Cell Disease

Al-Anazi KA\*

Consultant Hemato-Oncologist, Section of Adult Hematology and Oncology, Department of Medicine, College of Medicine and King Khalid University Hospital, King Saud University

### Abstract

Sickle cell anemia is associated with several systemic complications and life-threatening crises. The use of hydroxyurea, which increases hemoglobin F level in patients with sickle cell disease, is associated with reduction in disease severity, diminution in the frequency of veno-occlusive crises and other complications as well as improvement in the quality of life. There are specific indications for the use of hydroxyurea in patients with sickle cell disease and there are certain tools to monitor its effectiveness.

Hydroxyurea has been shown to be a safe medication even on long-term administration. However, close monitoring and regular follow up of patients from clinical and laboratory sides are essential to ensure early discovery of toxicity, adherence to the prescribed treatment in addition to continued and long-lasting response to therapy. In this literature review: the complications of sickle cell anemia as well as the available therapeutic modalities will be highlighted and the role of hydroxyurea in the management of patients with the disease will be discussed thoroughly.

**Keywords:** Sickle cell disease; Sickle cell crises; Hydroxyurea; Hemoglobin S; Hemoglobin F; Adherence to treatment

### Introduction

Sickle cell disease (SCD) is a very devastating medical condition caused by an autosomal recessive inherited hemoglobinopathy. This genetic defect is characterized by production of an abnormal hemoglobin (Hb) called Hb S which is poorly soluble and becomes polymerized when deoxygenated [1]. The pathophysiological hallmark of SCD is intracellular polymerization of Hb S upon deoxygenation [2]. Hb S results from point mutation in  $\beta$ -globin chain of human Hb (Glu 6 Val) where the sixth amino acid, glutamine, is substituted by valine [1,3-5]. This genetic mutation causes red blood cells (RBCs) to acquire a sickle shape under conditions of hypoxia resulting in a wide range of complications such as veno-occlusive crises, hemolytic episodes, stroke, acute chest syndrome and susceptibility to various infections [4].

Worldwide, Hb S is the commonest pathological Hb mutation and is one of the most common genetic causes of illness and death [5,6]. Recently, SCD has been recognized as a global health problem by the United Nations (UN) and the world health organization (WHO) [6]. Currently, the global burden of SCD is increasing and is expected to increase further in the coming decades [6,7]. SCD occurs throughout sub-Saharan Africa and in small pockets in the: Middle East, Indian subcontinent and Mediterranean and Caribbean regions [8,9]. Worldwide, approximately 300,000 to 400,000 children are born every year with SCD and up to 90% of these births occur in low-or middle-income countries [5,8].

### Complications of SCD and General Outline of Management

Patients with SCD are at risk of developing a number of life-threatening crises and the disease may be complicated by adverse consequences that may involve any body organ (Table 1) [1-4,10,11]. A study that was performed in both the United Kingdom (UK) and the United States of America (USA) which included 632 patients showed that the risk factors for death in patients with SCD include: age > 47 years, male gender, chronic blood transfusions, WHO class III-IV, and elevated hemolytic markers, serum ferritin and serum creatinine level [12]. Care of patients with SCD is a difficult task that requires a multidisciplinary approach [1,3,11,13]. Management of patients with SCD can be divided into: (1) general measures that can serve as long-term care as well as

crisis intervention, (2) specific and targeted therapies that are gaining particular interest and are aimed at prevention of complications and management of these complications once they are encountered, (3) potentially curative therapies and (4) preventive measures (Table 2) [1,3,11,14-21].

### Targeted Therapies in the Management of SCD

There are several classes of targeted and specific therapies that have been utilized in the treatment of patients with SCD. Examples of these drug categories include: drugs that increase Hb F production, Gardos channel inhibitors, anti-oxidants and vascular tone targeting agents (Table 2) [3,22-25]. Amongst the targeted therapies, agents that increase production of Hb F have received especial attention. Hydroxyurea is the most commonly studied and used agent [3,22,23]. Gardos channel inhibitors, such as ICA-17043, have been reported to improve RBC survival but only minimal improvement on pain episodes has been achieved. These drugs modulate the transport systems that are involved in cellular dehydration [1,3]. Because many factors in sickle-cell-induced ischemic injury are regulated by nitrous oxide (NO), the role of NO has been explored in patients with SCA. While pilot studies treating individuals with NO initially showed beneficial effects, larger studies did not show similar benefits [3].

### Hydroxyurea in the Treatment of SCD

Hydroxyurea was initially synthesized in Germany in 1869. It is an oral chemotherapeutic agent that has traditionally been used in the treatment of: chronic myeloproliferative neoplasms, certain types of leukemia, melanoma and ovarian cancer [26]. Hydroxyurea was first tested in SCD in 1984. In the USA, the drug was approved

**\*Corresponding author:** Khalid Ahmed Al-Anazi, Consultant Hemato-Oncologist, Section of Adult Hematology and Oncology, Department of Medicine, College of Medicine and King Khalid University Hospital, King Saud University, P.O. Box: 2925, Riyadh 11461, Saudi Arabia, Tel: 966- 011- 4671546; Fax: 966 - 011- 4671546; E-mail: [kaa\\_alanazi@yahoo.com](mailto:kaa_alanazi@yahoo.com)

**Received** December 15, 2014; **Accepted** December 25, 2014; **Published** January 05, 2015

**Citation:** Al-Anazi KA (2015) Hydroxyurea Therapy in Patients with Sickle Cell Disease. Transl Med 5: 145. doi:10.4172/2161-1025.1000145

**Copyright:** © 2015 Al-Anazi KA. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited



Body organ / system	Specific adverse effects
Hematopoietic system	* Variable degrees of myelosuppression, usually reversible: - mild thrombocytopenia - severe anemia - mild to moderate neutropenia * Megaloblastic erythropoiesis
Gastrointestinal tract	- Nausea - Vomiting - Diarrhea - Constipation
Liver	* Hyperbilirubinemia * Elevated liver enzymes
Skin, hair and nails	** Various skin eruptions: - Erythematous rash - Facial erythema - Maculopapular rash - Hyperpigmentation ** Skin ulcerations ** Hair loss ** Nail changes
Fertility in males	* Reduced sperm count - semen impairment appears to be proportional to the duration of hydroxyurea therapy. * Reduced sperm motility * Abnormal sperm morphology -semen impairment is usually irreversible in the majority of cases.
Miscellaneous side effects	* Renal tubular dysfunction ** Headache ** Fever ** Weight loss * Pulmonary fibrosis - rarely * Parvovirus-B19 infection ** Few reports of leukemia in humans.
<b>Adverse effects in animal studies</b>	** Carcinogenesis ** Teratogenesis ** Exacerbation of SCD - induced hypogonadism and development of gonadal failure ultimately.

**Table 3:** Adverse effects of hydroxyurea in humans and animal studies.

Indications	Details and examples
(1) Acute veno-occlusive complications	- Recurrent painful events - Dactylitis - Acute chest syndrome - Frequent hospitalizations
(2) Laboratory markers of severity	- Low hemoglobin - High white blood cell count - Low hemoglobin F level - High lactic dehydrogenase level
(3) Organ dysfunction	- Renal disease; proteinuria - Pulmonary disease; hypoxemia - Brain: * Elevated TCD velocities * Silent MRI or MRA changes * Stroke prophylaxis
(4) Miscellaneous	- Poor growth parameters - Patient or family request - Sibling on treatment

- TCD: transcranial doppler  
- MRI: magnetic resonance imaging  
- MRA: magnetic resonance angiography

**Table 4:** Potential clinical indications for using hydroxyurea in the treatment of patients with sickle cell disease

The use of hydroxyurea has been associated with adverse effects that can involve any body organ, but these side effects are not usually severe (Table 3)[21,30,38-42]. The effect of hydroxyurea on renal function is controversial. One study showed that the drug decreases glomerular hyperfiltration in children with SCD, while two other studies showed that the use of hydroxyurea in patients with SCD is associated with not only preservation but also improvement in renal concentrating ability and even reduction in renal enlargement [43-45]. A three year follow up of patients with SCD treated with hydroxyurea showed no major short or medium-term toxicity and a 9 year follow up study showed that no serious adverse effects were encountered on long-term use of the drug [39,46]. Due to the safety and efficacy of hydroxyurea in very young children, it is recommended to consider hydroxyurea therapy for all children with SCD regardless the presence or absence of clinical symptoms [47-49].

The potential indications for the use of hydroxyurea in infants, children and adults with SCD are included in Table 4 [22,50-53]. Several studies on the use of hydroxyurea in patients with SCD have shown the following beneficial effects: (1) increased production of Hb F with a concomitant reduction in the intracellular concentration of Hb S that can lead to an increase in total Hb concentration and decreased hemolysis with the release of free Hb, (2) reduction in white blood cell (WBC) count and reduction in the expression of cell

adhesion molecules that contribute to diminution of veno-occlusive crises and other complications, (3) reduction in the rate of transfusion requirements, (4) reduction in the number of visits to the emergency room and in the frequency of hospital admissions, (5) reduction in the rate of infectious complications, (6) significant reductions in the costs of medical care, and (7) reduction in the severity of SCD, decreased mortality, improved survival and improvement in the quality of life [22,30,46,50,51,54-56]. Hence, the use of hydroxyurea in patients with SCD is cost-effective [60,61]. A prospective, two arm, single center study on the effect of long-term administration of hydroxyurea on morbidity and mortality in adults with SCD that included 330 patients showed: the probability of 10 year survival was 86% and 65% for hydroxyurea and non-hydroxyurea treated patients respectively ( $p=0.001$ ) although hydroxyurea patients had more severe forms of SCD. Multivariate analysis showed that Hb F values at baseline and percentage change in lactic dehydrogenase (LDH) between baseline and 6 months were independent predictors of survival in the hydroxyurea group of patients [56]. Therefore, hydroxyurea treatment is not only beneficial in patients with SCD, but its long-term use can also modify the natural history of the disease [53]. However, the use of hydroxyurea in children with SCD is significantly associated with: (1) sickle genotype, (2) better parental knowledge about the major therapeutic effects of the drug, and (3) institution of this therapy by hematologists and provision of the drug to symptomatic patients [23].

Blood parameters	Details
Neutrophils	Absolute neutrophil count <math> < 0.1 \times 10^9 / L </math>
Hemoglobin	<math> < 7.0 \text{ g/dL}</math> with low reticulocyte count [absolute reticulocyte count >math> > 100 \times 10^9 / L </math>] or Decrease in hemoglobin by > 20% from previous level
Reticulocytes	<math> < 80 \times 10^9 / L </math> unless hemoglobin concentration is > 8.0 gram / dL
Platelets	Platelet count <math> < 80 \times 10^9 / L </math>

- L: liter  
 - dL: decilite

**Table 5:** Hematological toxicity thresholds requiring hydroxyurea dose modification.

Predictors of response to hydroxyurea therapy include: (1) decrease in the number of veno-occlusive crises and reduction in the complications of SCD, (2) increase in normal Hb level, Hb F level, mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean reticulocyte volume and percentage of F-cells, and (3) increase in the urinary level of hydroxyurea as measured by mass spectrometry [62-68]. Other parameters that can predict the response to hydroxyurea therapy include: duration of treatment with hydroxyurea, certain haplotypes such as Senegal haplotype, gender and weight of patient, WBC count, platelet count and reticulocyte count [68]. Sometimes, the dose of hydroxyurea may need to be modified due to the evolution of certain adverse effects that may be encountered on the long-term use of the drug. Hematological toxicity thresholds that require modification of the dose of hydroxyurea in patients with SCD are included in Table 5 [22,52].

Adherence to a medication is defined as the extent to which patients take medications as prescribed by their health care providers. Adherence to drug therapy can be measured by direct and indirect methods, but none of them is considered the gold standard. Direct methods include: directly observed therapy (DOT) and measurement of drug levels or biologic markers in blood. Indirect methods include: pill counts, pharmacy refills, electronic drug monitoring, patient diaries and patient self-reports the form of questionnaires [69]. The modified version of Morisky medication adherence scale (MMAS) is relatively simple, practical to use in clinical settings and is preferred to interviews. The eight items included in the scale aid in identifying patients with adherence problems and can be utilized to monitor adherence over the course of drug therapy [70-73]. Adherence to hydroxyurea can be measured by: (1) pharmacy refills, (2) MMAS, (3) visual analogue scale, (4) medical provider report, (5) clinic visits and (6) electronic DOT [74,75]. Pharmacy refills and MMAS may be helpful in identifying children at risk of poor response due to non-adherence and children with good adherence but having poor response due to biologic factors such as pharmacodynamics [74]. Studies have shown that close adherence to prescribed hydroxyurea therapy is necessary to maximize the efficacy of the drug and that adherence to treatment improves health-related quality of life [74,76-78]. In a larger retrospective study which was conducted in North Carolina in the USA and that included 312 patients with SCD treated with hydroxyurea, the following results were obtained: (1) adherence to treatment was generally suboptimal, as nearly two thirds of the study population were classified as non-adherent to therapy, and (2) adherence to hydroxyurea was associated with: reduced risk of hospitalization, reduction in the number of visits to the emergency room, reduction in the frequency of veno-occlusive crises and reduction in health care costs [77]. In a multicenter, double-blinded, placebo-controlled study on the efficacy of hydroxyurea therapy in SCA patients which included 150 patients treated with hydroxyurea and that was performed in USA in 1995, the results of follow up for 2 years showed: (1) half of the hydroxyurea-assigned patients had long-term increment in Hb F, and (2) bone marrow reserve or ability to withstand hydroxyurea therapy was important for sustained Hb F

increase during treatment with hydroxyurea [62]. Extended follow up of patients for 17.5 years showed significant reduction in mortality and safety of the drug on long-term use [55].

Barriers to adherence to hydroxyurea treatment include: high frequency and length of clinic visits, disruption of school and work and lack of resources [78]. Provider-related barriers to prescribing hydroxyurea include: patient adherence to the prescribed medication, patient adherence to blood tests and lack of contraception in females [79]. Facilitators of adherence include: health benefits of therapy, social support systems, medication reminders and positive clinic experiences [78]. In addition to health education, the following methods have been shown to improve adherence to treatment: (1) multimedia communication for hospitalized patients with their teachers and work colleagues, (2) integration of interactive web-based technology to assess adherence to therapy, (3) electronic DOT, and (4) coaching very young children for swallowing hydroxyurea capsules [75,78,80-82].

Over the last 30 years, substantial experience has accumulated regarding the safety and efficacy of hydroxyurea in patients with SCD and strong evidence supports its efficacy in reducing the frequency of veno-occlusive crises, acute chest syndrome, hospitalizations and blood transfusions in patients with SCD [29,50,51,53,54]. Thus, hydroxyurea is considered an important advance in the treatment of patients with SCD as it is the only widely used drug that modifies the pathogenesis of the disease [29,51-53,55]. Although questions remain regarding its long-term risks and benefits, current evidence suggests that many young patients with SCD should receive hydroxyurea [29,50,51,53,54].

### Curative And Potentially Curative Therapies

Potentially curative therapies in patients with SCD include: gene replacement therapy and hematopoietic stem cell transplantation (HSCT) [1,3,16,18-21]. Gene replacement therapy, using either pluripotent stem cells or autologous gene correction in stem cell designs, has been studied in animal models with promising results [17,18]. Induced pluripotent stem cells (iPSCs) have the potential to correct the Glu-6-Val mutation in the abnormal hemoglobin (Hb S). However, safety issues are of a concern [3,16-19]. The only curative therapy for SCD is allogeneic HSCT. Traditionally this procedure has been limited to children below 16 years of age with severe pre-existing complications [1,19-21]. Use of HLA-matched sibling donor myeloablative allogeneic HSCT has been associated with: overall survival >90% and cure rate up to 85% [19-21]. However, HSCT particularly myeloablative allogeneic transplantation carries significant morbidity and mortality [19-21,83]. In sibling allografts, stem cells can be obtained from a normal donor or a sickle cell trait. Sibling allogeneic HSCT may be available for 30% of patients, while matched unrelated donor (MUD) allografts can be available for 60% of patients [20,21]. Recently, reduced-intensity conditioning therapy has been introduced in HSCT for patients with SCD and this form of HSCT allows older patients and those with organ dysfunction to be transplanted. Alternative donors such as unrelated cord blood HSCT as well as related haploidentical bone marrow and



peripheral blood HSCT have also been successfully used [19-21,83-85]. Interestingly, the use of hydroxyurea in patients with severe SCD prior to allogeneic HSCT does not have a negative impact on the outcome of HSCT as it seems to be associated with lower incidence of rejection and absent engraftment [86].

## Conclusions and Recommendations

Hydroxyurea treatment is effective in reducing the complications of SCD in infants, children and adults and it has significantly improved the quality of life in patients receiving it. However, adherence to the prescribed medication is essential to achieve the expected response. Close follow up of patients and regular monitoring of their clinical and laboratory parameters is vital to predict the response to hydroxyurea treatment. Adopting a multidisciplinary approach and incorporation of continued health education are important to guarantee regular intake of the drug in order to prevent complications of SCD before they evolve.

## References

- Kaur M, Dangi CBS, Singh M (2013) An overview of sickle cell disease profile. *Asian J Pharmaceut Clin Res* 6 : 25-37.
- Schnog JB1, Duits AJ, Muskiet FA, ten Cate H, Rojer RA, et al. (2004) Sickle cell disease; a general overview. *Neth J Med* 62: 364-374.
- Bender MA, Hobbs W. Sickle cell disease. *Gene Reviews* [Internet] - NCBI Bookshelf 2012. Pagon RA, Adam MP, Ardinger HH, et al, editors. Seattle (WA): University of Washington, Seattle; 1993-2014. Initial posting: September 15, 2003. Last revision: May 17, 2012. Pages: 1-67.
- Driss A, Asare KO, Hibbert JM, Gee BE, Adamkiewicz TV, et al. (2009) Sickle Cell Disease in the Post Genomic Era: A Monogenic Disease with a Polygenic Phenotype. *Genomics Insights* 2009: 23-48.
- Makani J, Ofori-Acquah SF, Nnodu O3, Wonkam A4, Ohene-Frempong K5 (2013) Sickle cell disease: new opportunities and challenges in Africa. *ScientificWorldJournal* 2013: 193252.
- Piel FB, Patil AP, Howes RE, Nyangiri OA, Gething PW, et al. (2013) Global epidemiology of sickle haemoglobin in neonates: a contemporary geostatistical model-based map and population estimates. *Lancet* 381: 142-151.
- Piel FB, Hay SI, Gupta S, Weatherall DJ, Williams TN (2013) Global burden of sickle cell anaemia in children under five, 2010-2050: modelling based on demographics, excess mortality, and interventions. *PLoS Med* 10: e1001484.
- Williams TN, Weatherall DJ (2012) World distribution, population genetics, and health burden of the hemoglobinopathies. *Cold Spring Harb Perspect Med* 2: a011692.
- El-Hazmi MA, Al-Hazmi AM, Warsy AS (2011) Sickle cell disease in Middle East Arab countries. *Indian J Med Res* 134: 597-610.
- Vichinsky EP (2014) Overview of the clinical manifestations of sickle cell disease. *UpToDate*. Schrier SL and Tirnauer JS.(Eds) Topic 7119: 1-14.
- Ballas SK, Kesen MR, Goldberg MF, Luty GA, Dampier C, et al. (2012) Beyond the definitions of the phenotypic complications of sickle cell disease: an update on management. *ScientificWorldJournal* 2012: 949535.
- Gladwin MT, Barst RJ, Gibbs JS, Hildesheim M, Sachdev V, et al. (2014) Risk factors for death in 632 patients with sickle cell disease in the United States and United Kingdom. *PLoS One* 9: e99489.
- Apanah S, Rizzolo D (2013) Sickle cell disease: taking a multidisciplinary approach. *JAAAPA* 26: 28-33.
- Field JJ, Vichinsky EP, DeBaun MR (2014) Overview of the management and prognosis of sickle cell disease. *UpToDate*. Schrier SL, Mahoney Jr DH, Tirnauer JS. (Eds).
- Manwani D, Frenette PS (2013) Vaso-occlusion in sickle cell disease: pathophysiology and novel targeted therapies. *Blood* 122: 3892-3898.
- Frenette PS, Atweh GF (2007) Sickle cell disease: old discoveries, new concepts, and future promise. *J Clin Invest* 117: 850-858.
- Hanna J, Wernig M, Markoulaki S, Sun CW, Meissner A, et al. (2007) Treatment of sickle cell anemia mouse model with iPS cells generated from autologous skin. *Science* 318: 1920-1923.
- Townes TM (2008) Gene replacement therapy for sickle cell disease and other blood disorders. *Hematology Am Soc Hematol Educ Program* .
- Freed J, Talano J, Small T, Ricci A, Cairo MS (2012) Allogeneic cellular and autologous stem cell therapy for sickle cell disease: 'whom, when and how'. *Bone Marrow Transplant* 47: 1489-1498.
- Shenoy S (2011) Hematopoietic stem cell transplantation for sickle cell disease: current practice and emerging trends. *Hematology Am Soc Hematol Educ Program* 2011: 273-279.
- Gluckman E (2013) Allogeneic transplantation strategies including haploidentical transplantation in sickle cell disease. *Hematology Am Soc Hematol Educ Program* 370-376.
- Strouse JJ, Heeney MM (2012) Hydroxyurea for the treatment of sickle cell disease: efficacy, barriers, toxicity, and management in children. *Pediatr Blood Cancer* 59: 365-371.
- Oyeku SO, Driscoll MC, Cohen HW, Trachtman R, Pashankar F, et al. (2013) Parental and other factors associated with hydroxyurea use for pediatric sickle cell disease. *Pediatr Blood Cancer* 60: 653-658.
- Eridani S, Mosca A (2011) Fetal hemoglobin reactivation and cell engineering in the treatment of sickle cell anemia. *J Blood Med* 2: 23-30.
- Trompeter S, Roberts I (2009) Haemoglobin F modulation in childhood sickle cell disease. *Br J Haematol* 144: 308-316.
- Brawley OW, Cornelius LJ, Edwards LR, Gamble VN, Green BL, et al. (2008) National Institutes of Health Consensus Development Conference Statement: hydroxyurea treatment for sickle cell disease. *Ann Intern Med* 148: 932-938.
- Ware RE (2010) How I use hydroxyurea to treat young patients with sickle cell anemia. *Blood* 115: 5300-5311.
- Mulaku M, Opiyo N, Karumbi J, Kitonyi G, Thoithi G, et al. (2013) Evidence review of hydroxyurea for the prevention of sickle cell complications in low-income countries. *Arch Dis Child* 98: 908-914.
- Segal JB, Strouse JJ, Beach MC, Haywood C, Witkop C, et al. (2008) Hydroxyurea for the treatment of sickle cell disease. *Evidence Report Technology Assessment Number 165*. AHRQ Publication No. 08. E007.
- Odievre M-H, Benkerrou M, Belloy M, Elion J (2008) Hydroxyurea treatment in sickle cell children. *Paed Perinat Drug Ther* 8: 158-170.
- Green NS1, Barral S2 (2014) Emerging science of hydroxyurea therapy for pediatric sickle cell disease. *Pediatr Res* 75: 196-204.
- Sheehan VA, Luo Z, Flanagan JM, Howard TA, Thompson BW, et al. (2013) Genetic modifiers of sickle cell anemia in the BABY HUG cohort: influence on laboratory and clinical phenotypes. *Am J Hematol* 88: 571-576.
- Kumkhaek C, Taylor JG 6th, Zhu J, Hoppe C, Kato GJ, et al. (2008) Fetal haemoglobin response to hydroxycarbamide treatment and sar1a promoter polymorphisms in sickle cell anaemia. *Br J Haematol* 141: 254-259.
- McGann PT, Flanagan JM, Howard TA, Dertinger SD, He J, et al. (2012) Genotoxicity associated with hydroxyurea exposure in infants with sickle cell anemia: results from the BABY-HUG Phase III Clinical Trial. *Pediatr Blood Cancer* 59: 254-257.
- Khayat AS, Antunes LM, Guimarães AC, Bahia MO, Lemos JA, et al. (2006) Cytotoxic and genotoxic monitoring of sickle cell anaemia patients treated with hydroxyurea. *Clin Exp Med* 6: 33-37.
- Lederman HM, Connolly MA, Kalpathi R, Ware RE, Wang WC, et al. (2014) Immunologic effects of hydroxyurea in sickle cell anemia. *Pediatrics* 134: 686-695.
- Benito JM, López M, Lozano S, Ballesteros C, González-Lahoz J, et al. (2007) Hydroxyurea exerts an anti-proliferative effect on T cells but has no direct impact on cellular activation. *Clin Exp Immunol* 149: 171-177.
- Ballas SK, Singh P, Adams-Graves P, Wordell CJ (2013) Idiosyncratic side effects of hydroxyurea in patients with sickle cell anemia. *J Blood Disorders Transf* 4:5.
- de Montalembert M, Bégue P, Bernaudin F, Thuret I, Bachir D, et al. (1999) Preliminary report of a toxicity study of hydroxyurea in sickle cell disease. *French Study Group on Sickle Cell Disease. Arch Dis Child* 81: 437-439.
- Jones KM, Niaz MS, Brooks CM, Roberson SI, Aguinaga MP, et al. (2009) Adverse effects of a clinically relevant dose of hydroxyurea used for the treatment of sickle cell disease on male fertility endpoints. *Int J Environ Res Public Health* 6: 1124-1144.

41. Lukusa AK, Vermeylen C, Vanabelle B, Curaba M, Brichard B, et al. (2009) Bone marrow transplantation or hydroxyurea for sickle cell anemia: long-term effects on semen variables and hormone profiles. *Pediatr Hematol Oncol* 26: 186-194.
42. Grigg A (2007) Effect of hydroxyurea on sperm count, motility and morphology in adult men with sickle cell or myeloproliferative disease. *Intern Med J* 37: 190-192.
43. Aygun B, Mortier NA, Smeltzer MP, Shulkin BL, Hankins JS, et al. (2013) Hydroxyurea treatment decreases glomerular hyperfiltration in children with sickle cell anemia. *Am J Hematol* 88: 116-119.
44. Miller ST, Wang WC, Iyer R, Rana S, Lane P, et al. (2010) Urine concentrating ability in infants with sickle cell disease: baseline data from the phase III trial of hydroxyurea (BABY HUG). *Pediatr Blood Cancer* 54: 265-268.
45. Alvarez O, Miller ST, Wang WC, Luo Z, McCarville B, et al and the BABY HUG Investigators (2012) Effect of hydroxyurea treatment on renal function parameters: results from the multi-center placebo-controlled BABY HUG clinical trial for infants with sickle cell anemia. *Pediatr Blood Cancer* 59 (4): 668-674.
46. Steinberg MH, Barton F, Castro O, Pegelow CH, Ballas SK, et al. (2003) Effect of hydroxyurea on mortality and morbidity in adult sickle cell anemia: risks and benefits up to 9 years of treatment. *JAMA* 289: 1645-1651.
47. Thornburg CD, Files BA, Luo Z, Miller ST, Kalpathi R, et al. (2012) Impact of hydroxyurea on clinical events in the BABY HUG trial. *Blood* 120: 4304-4310.
48. Wang WC, Ware RE, Miller ST, Iyer RV, Casella JF, et al. (2011) Hydroxycarbamide in very young children with sickle-cell anaemia: a multicentre, randomised, controlled trial (BABY HUG). *Lancet* 377: 1663-1672.
49. Zimmerman SA, Schultz WH, Davis JS, Pickens CV, Mortier NA, et al. (2004) Sustained long-term hematologic efficacy of hydroxyurea at maximum tolerated dose in children with sickle cell disease. *Blood* 103: 2039-2045.
50. Rodgers GP (2014) Hydroxyurea and other disease-modifying therapies in sickle cell disease. *UpToDate*. Schrier SL, Mahoney DH and Tirnauer JS (Eds).
51. Yawn BP, Buchanan GR, Afenyi-Annan AN, Ballas SK, Hassell KL, et al. (2014) Management of sickle cell disease: summary of the 2014 evidence-based report by expert panel members. *JAMA* 312: 1033-1048.
52. Heeney MM1, Ware RE (2010) Hydroxyurea for children with sickle cell disease. *Hematol Oncol Clin North Am* 24: 199-214.
53. Ware RE, Aygun B (2009) Advances in the use of hydroxyurea. *Hematology Am Soc Hematol Educ Program*.
54. Silva-Pinto AC, Angulo IL, Brunetta DM, Neves FI, Bassi SC, et al. (2013) Clinical and hematological effects of hydroxyurea therapy in sickle cell patients: a single-center experience in Brazil. *Sao Paulo Med J* 131: 238-243.
55. Steinberg MH, McCarthy WF, Castro O, Ballas SK, Armstrong FD, et al. (2010) The risks and benefits of long-term use of hydroxyurea in sickle cell anemia: A 17.5 year follow-up. *Am J Hematol* 85: 403-408.
56. Voskaridou E, Christoulas D, Billalis A, Plata E, Varvagiannis K, et al (2010) The effect of prolonged administration of hydroxyurea on morbidity and mortality in adult patients with sickle cell syndromes: results of a 17-year, single-center trial (LaSHS). *Blood* 115 : 2354-2363.
57. Thornburg CD, Calatroni A, Panepinto JA (2011) Differences in health-related quality of life in children with sickle cell disease receiving hydroxyurea. *J Pediatr Hematol Oncol* 33: 251-254.
58. Nwenyi E, Leafman J, Mathieson K, Ezeobah N (2014) Differences in quality of life between pediatric sickle cell patients who used hydroxyurea and those who did not. *Int J Health Care Qual Assur* 27: 468-481.
59. Nottage KA, Hankins JS, Smeltzer M, Mzayek F, Wang WC, et al. (2013) Hydroxyurea use and hospitalization trends in a comprehensive pediatric sickle cell program. *PLoS One* 8: e72077.
60. Wang WC, Oyeku SO, Luo Z, Boulet SL, Miller ST, et al. (2013) Hydroxyurea is associated with lower costs of care of young children with sickle cell anemia. *Pediatrics* 132: 677-683.
61. Stallworth JR, Jerrell JM, Tripathi A (2010) Cost-effectiveness of hydroxyurea in reducing the frequency of pain episodes and hospitalization in pediatric sickle cell disease. *Am J Hematol* 85: 795-797.
62. Steinberg MH, Lu ZH, Barton FB, Terrin ML, Charache S, et al. (1997) Fetal hemoglobin in sickle cell anemia: determinants of response to hydroxyurea. Multicenter Study of Hydroxyurea. *Blood* 89: 1078-1088.
63. Borba R, Lima CS, Grotto HZ (2003) Reticulocyte parameters and hemoglobin F production in sickle cell disease patients undergoing hydroxyurea therapy. *J Clin Lab Anal* 17: 66-72.
64. Ware RE, Eggleston B, Redding-Lallinger R, Wang WC, Smith-Whitley K, et al. (2002) Predictors of fetal hemoglobin response in children with sickle cell anemia receiving hydroxyurea therapy. *Blood* 99: 10-14.
65. Ma Q, Wyszynski DF, Farrell JJ, Kutlar A, Farrer LA, et al. (2007) Fetal hemoglobin in sickle cell anemia: genetic determinants of response to hydroxyurea. *Pharmacogenomics J* 7: 386-394.
66. Mach Queiroz AM, de Castro Lobo CL, do Nascimento EM, de Braganca Pereira B, Bonini-Domingo CR, et al. (2013) The mean corpuscular volume and hydroxyurea in Brazilian patients with sickle cell anemia: a surrogate marker of compliance. *J Blood Disorders Transf* 4: 157.
67. Dalton RN, Turner C, Dick M, Height SE, Awogbade M, et al. (2005) The measurement of urinary hydroxyurea in sickle cell anaemia. *Br J Haematol* 130: 138-144.
68. Valafar H, Valafar F, Darvill A, Albersheim P, Kutlar A, et al. (2000) Predicting the effectiveness of hydroxyurea in individual sickle cell anemia patients. *Artif Intell Med* 18: 133-148.
69. Osterberg L, Blaschke T (2005) Adherence to medication. *N Engl J Med* 353: 487-497.
70. Morisky DE, Green LW, Levine DM (1986) Concurrent and predictive validity of a self-reported measure of medication adherence. *Med Care* 24: 67-74.
71. Morisky DE, Ang A, Krousel-Wood M, Ward HJ (2008) Predictive validity of a medication adherence measure in an outpatient setting. *J Clin Hypertens (Greenwich)* 10: 348-354.
72. Al-Qazaz HKh, Hassali MA, Shafie AA, Sulaiman SA, Sundram S, et al. (2010) The eight-item Morisky Medication Adherence Scale MMAS: translation and validation of the Malaysian version. *Diabetes Res Clin Pract* 90: 216-221.
73. Garber MC, Nau DP, Erickson SR, Aikens JE, Lawrence JB (2004) The concordance of self-report with other measures of medication adherence: a summary of the literature. *Med Care* 42: 649-652.
74. Thornburg CD, Calatroni A, Telen M, Kemper AR (2010) Adherence to hydroxyurea therapy in children with sickle cell anemia. *J Pediatr* 156: 415-419.
75. Creary SE, Gladwin MT, Byrne M, Hildesheim M, Krishnamurti L (2014) A pilot study of electronic directly observed therapy to improve hydroxyurea adherence in pediatric patients with sickle-cell disease. *Pediatr Blood Cancer* 61: 1068-1073.
76. Al Jaouni SK, Al Muhayawi MS, Halawa TF, Al Mehayawi MS (2013) Treatment adherence and quality of life outcomes in patients with sickle cell disease. *Saudi Med J* 34: 261-265.
77. Candrilli SD, O'Brien SH, Ware RE, Nahata MC, Seiber EE, et al. (2011) Hydroxyurea adherence and associated outcomes among Medicaid enrollees with sickle cell disease. *Am J Hematol* 86: 273-277.
78. Boateng L, Puffer E, Allen J, Bonner M, Thornburg CD (2013) Adherence to secondary stroke prevention in children with sickle cell anemia: family and child perspectives. *J Blood Disorders Transf* 4 : 148.
79. Brandow AM, Jirovec DL, Panepinto JA (2010) Hydroxyurea in children with sickle cell disease: practice patterns and barriers to utilization. *Am J Hematol* 85: 611-613.
80. Grove J, Grove O, Michie C (2013) The real effects of sickle cell disease on children and adolescents. *West London Med J* 6 : 11-17.
81. Crosby LE, Barach I, McGrady ME, Kalinyak KA, Eastin AR, et al. (2012) Integrating interactive web-based technology to assess adherence and clinical outcomes in pediatric sickle cell disease. *Anemia* 2012: 492428.
82. Bekele E, Thornburg CD, Brandow AM, Sharma M, Smaldone AM, et al. (2014) Do difficulties in swallowing medication impede the use of hydroxyurea in children? *Pediatr Blood Cancer* 61: 1536-1539.
83. Hsieh MM, Fitzhugh CD1, Weitzel RP1, Link ME1, Coles WA1, et al. (2014) Nonmyeloablative HLA-matched sibling allogeneic hematopoietic stem cell transplantation for severe sickle cell phenotype. *JAMA* 312: 48-56.
84. King AA, DiPersio JF2 (2014) Reconsideration of age as a contraindication for curative therapy of sickle cell disease. *JAMA* 312: 33-34.
85. Talano JA, Cairo MS (2014) Hematopoietic stem cell transplantation for sickle cell disease: state of the science. *Eur J Haematol*.
86. Brachet C, Azzi N, Demulder A, Devalck C, Gourdin A, et al. (2004) Hydroxyurea treatment for sickle cell disease: impact on haematopoietic stem cell transplantation's outcome. *Bone Marrow Transplant* 33: 799-803.