

## Evaluation the Efficacy of *Phytolacca dodecandra*, *Buddleia polystachya* and *Vernonia amygdalina* Leaf Extract against German Cockroach (*Blattella germanica*) under Uncontrolled Condition

Weldesent Beze Kassie \*

Department of Food Science, Central Ethiopian Environment and Forest Research Center, Addis Ababa, Ethiopia

### ABSTRACT

The present study was conducted to evaluate the efficacy of *Phytolacca dodecandra*, *Buddleia polystachya* and *Vernonia amygdalina* leaf extract against the adult German cockroach, *Blattella germanica* under uncontrolled condition by comparing control measures. The research consists of three botanical leaf extracts were prepared by three concentration levels (5 g/100 ml, 10 g/100 ml and 15 g/100 ml of water), a positive control (deltamethrin) and negative control (water), were laid out in a randomized complete design with three replications. Total of 330 adult cockroaches were tested. Mortality of cockroaches was recorded after 24, 48 and 72 hours of exposure. Data were analyzed by using SPSS software package version 20.0. The results indicated that all the botanical extract concentrations were significant for cockroaches mortality. The mean mortality count of adult *B. germanica* was the highest (100) in 15 g/100 ml concentration of *P. dodecandra* and positive control and the lowest (no mortality) in the negative control was recorded in 24 hours exposure time. Hundred percent (100%) of mortality were recorded by *P. dodecandra* at 5 g/100 ml after 72 hours, 10 g/100 ml after 48 hours 15 g/100 ml after 24 hours and by *B. polystachya* and *V. amygdalina* at 15 g/100 ml after 48 hours. These results were no significant differences with positive control. But, there was significant difference in mortality in both increase in concentration levels and exposure time for all botanicals. Therefore, users use these easily prepared and available botanicals to control/eliminate cockroaches by formulating 15 g/100 ml of water and more concentration levels rather than using of synthetic pesticides to keep environmental pollution.

**Keywords:** *Blattella germanica*, botanical pesticides, efficacy, extract, mortality

### INTRODUCTION

The increased use of pesticides in the control of pests has prompted some concern over the adverse effects of these chemical substances on living organisms and their environment [1], thus researchers are now focusing attention to alternative method of controlling insect vectors [2]. Indigenous plants exist in nature, particularly in tropical areas, which are associated with some domestic uses [3]. These plants contain bioactive chemicals, which serve as suitable alternative biocontrol substances [4]. Thus repel, inhibit growth or kill pests [5]. Botanical pesticides prepared from their barks, leaves, roots, flowers, fruits, seeds, cloves, rhizomes and stems [6]. The plant

parts are dried and ground into fine powder and extracted with organic solvents that will maximize extraction of the targeted compounds. Then, extracts are concentrated, formulated and evaluated for efficacy under laboratory, controlled or field conditions [7]. Botanical pesticides are considered safe in pest control because they have low or none pesticide residue making them safe to the people, environment and ecosystem [8]; they affect insect pests in different ways depending on physiological characteristics of the insect species as well as the type of the insecticidal plants. The components of various botanical insecticidal can be classified into six groups namely; repellents, feeding deterrents/anti-feedants, toxicants, growth retardants, chemosterilants and attractants [9].

**Correspondence to:** Weldesenbet Beze Kassie, Department of Plant Science, Central Ethiopian Environment and Forest Research Center, Addis Ababa, Ethiopia, E-mail: weldeentomologyresearcher@gmail.com

**Received:** 08-Feb-2022, Manuscript No. JFMSH-20-5910; **Editor assigned:** 10-Feb-2022, PreQC No. JFMSH-20-5910 (PQ); **Reviewed:** 24-Feb-2022, QC No. JFMSH-20-5910; **Revised:** 02-Mar-2022, Manuscript No. JFMSH-20-5910 (R); **Published:** 09-Mar-2022, DOI: 10.35248/2476-2059-22.S3.004.

**Citation:** Kassie WB (2022) Evaluation the Efficacy of *Phytolacca dodecandra*, *Buddleia polystachya* and *Vernonia amygdalina* Leaf Extract against German Cockroach (*Blattella germanica*) under Uncontrolled Condition. Food Microbial Saf Hyg. S3:004.

**Copyright:** © 2022 Kassie WB. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Different botanicals are used to protect agricultural and household products as natural, cost effective and easily accessible from nearby source. In addition, due to their biodegradability unlike synthetic chemical pesticides they are environmental friendly and hence favorable to the environmental sustainability and community wellbeing [10]. They inhibit or disrupt insect feeding [9,11]; toxic cause death to insects [12]; showed deleterious effects on the growth and development of insects, reducing the weight of larva, pupa and adult stages and lengthening the development stages [11]; also reduce the survival rates of larvae and pupae as well as adult emergence [13]; causing temporary or permanent sterility of one or both of the sexes or preventing maturation of sexually functional adult stage [14,15].

Cockroaches are important insect pests of public health [16]; and the most common pests in many homes. Various studies around the world revealed that cockroaches living close to human dwellings were important carriers of etiologic agents belonging to all groups of potential pathogens and they cause of allergies and asthma. The insect is used as a key indicator of insanitary conditions [17,18]. The insect is as various agents vector, like a poliomyelitis virus, enter pathogenic bacteria, amebas cyst, eggs of worm, and the fungus *Aspergillus*. More or less 22 species of bacteria, virus, fungi and protozoans and 5 species of worms have been isolated from the body of the cockroach in the field [19]. The presence of cockroaches in homes can also cause psychological distress, elicit allergic reactions, and trigger asthma attacks as a result of the presence of allergens [20].

The German cockroach as a mechanical vector can easily transmit pathogens such as bacteria, viruses, fungi, protozoa, and parasites on their legs and bodies. It has a small body size, special nutritional habits, and specific behavior. Besides, it can readily adapt itself to the human environment and the causative agents of allergic diseases [21]. German cockroach found in most parts of the world. It is light yellowish brown and 10-15 mm in length, making it one of the smallest domestic cockroaches. The female usually carries the egg case until shortly before the young come out. It is considered as an important medical and economic pest because it has a shorter generation time and higher fecundity than the other cockroaches, which makes it difficult to control. The indiscriminate movement between filth and food make them potentially efficient vectors of human pathogens [22,20].

Conventional synthetic insecticides, such as pyrethroids, carbamates and organophosphates are very popularly used compounds for controlling cockroaches [23,24]. However, these insecticides have several adverse effects including acute or chronic toxicity to humans and animals. They can also have an adverse impact on the food chain of the ecosystem [25-27]. Furthermore, cockroaches can develop resistance to these insecticides. Due to these problems, an alternative approach is to search for effective botanical extracts, which are safe to humans and the environment [28]. The aim of this study is to evaluate the efficacy *P. dodecandra*, *B. polystachya* and *V. amygdalina* leaf aqua extracts to control adult *B. germanica* and to give information for users that use botanical extracts by easy

preparation method without cost and risk at places these cockroaches present.

## MATERIALS AND METHODS

### Locality

This experimental research study was done during the period of time February-April of the temperate season 2020, in the Yeka sub city, Addis Abeba, Ethiopia, which belongs to the average of temperature was 23.5°C (Table 1).

Scientific name	Family name	English name	Amharic name	Part used
<i>Phytolacca dodecandra</i>	Phytolaccaceae	Soap berry	Endod	Leaves
<i>Buddleia polystachya</i>	Scrophulariaceae	Butterfly bush	Anfar	Leaves
<i>Vernonia amygdalina</i>	Asteraceae	Bitter leaf	Grawa	Leaves

**Table 1:** Name of botanicals used in the experiment for against *Blattella germanica*.

### Collection and extraction of botanical Leaves

Fresh green *P. dodecandra*, *B. polystachya* and *V. amygdalina* leaves were collected from available areas and washed by water to remove dust and microorganisms that attached on their surfaces. Then put under shade for 15 days until they were dried completely. The dried leaves were and cut into pieces and ground by using an electrical grading machine until obtains very fine powder. The fine powder of each botanicals were weighed (5 g, 10 g, and 15 g) by sensitive weight balance. Then each balanced fine powder was mixed in 100 ml of water separately by 5 minutes shaking to made well solution. After staying 72 hours, each prepared solutions were filtered through mush cloth repeatedly until obtained 30 ml filtered extract for each treatments.

### Collection of cockroaches and study design

The experiment was laid out in a Randomized Complete Design (RCD) with three replications. Total more than 450 adult cockroaches were collected from 11 home rooms (their hidden places) by females by once night carefully without injure cockroaches. At morning, ten (10) cockroaches were counted and introduced into separately arranged 33 experimental units (containers). Total number of cockroaches for this efficiency test was 330. Botanicals extract concentrations and controls were sprayed in each experimental units these containing cockroaches. The number of died Cockroaches were observed, counted, recorded and removed after 24, 48 and 72 hours of spaying and death rate in (%) were calculated.

## Statistical analysis

Data collected were analyzed using one-way analysis of variance (ANOVA). Statistical significant differences of the treatment were determined using Turkey student test. All statistical analysis was carried out using SPSS version 20.0 statistical package.

## RESULTS AND DISCUSSION

Cockroaches are omnivorous and feed on human and non-human foods, such as household items, glue in books and furniture. They contaminate food by depositing saliva and faeces on it, and through the mechanical transfer of diseases, which can cause food poisoning and diarrhoea. They also have, and transmit; an unpleasant odour due to chemicals produced by abdominal glands and agitates allergies [29]. Plant extracts have been used worldwide as an alternative method to control pests unlike synthetic pesticide which have one active compound, and exhibit only one type of biological effect. Single plant derived compound may have more than one biological effect [30]. Plants are rich source of bioactive compounds that can be used as a suitable substitute to develop eco-friendly pest management strategies. In the present study, screening of the plant extracts along with the commercial botanicals showed that both possess bioactivity against *B. germanica*. The result of the present study showed that the mortality of *B. germanica* exposed to tested botanicals extract increased with increase in grams and exposure time. The mean mortality of adult *B. germanica* exposed to *P. dodecandra*, *B. polystachya* and *V. amygdalina* leaf extract is presented in Table 2.

Treatments	Concentration	Mean mortality $\pm$ SE over time (hours) after exposure			
		(w/v)	24 hours	48 hours	72 hours
<i>Phytolacca dodecandra</i>	5		66.00 $\pm$ 0.00	92.00 $\pm$ 0.00	100.00 $\pm$ 0.00
	10		91.00 $\pm$ 0.00	100.00 $\pm$ 0.00	100.00 $\pm$ 0.00
	15		100.00 $\pm$ 0.00	100.00 $\pm$ 0.00	100.00 $\pm$ 0.00
<i>Buddleia polystachya</i>	5		35.00 $\pm$ 0.00	66.00 $\pm$ 0.00	78.00 $\pm$ 0.00
	10		68.00 $\pm$ 0.00	90.00 $\pm$ 0.00	93.00 $\pm$ 0.00
	15		90.00 $\pm$ 0.00	100.00 $\pm$ 0.00	100.00 $\pm$ 0.00
<i>Vernonia amygdalina</i>	5		33.00 $\pm$ 0.00	42.00 $\pm$ 0.00	55.00 $\pm$ 0.00
	10		55.00 $\pm$ 0.00	64.00 $\pm$ 0.00	76.00 $\pm$ 0.00
	15		88.00 $\pm$ 0.00	100.00 $\pm$ 0.00	100.00 $\pm$ 0.00

Positive control	30 ml	100.00 $\pm$ 0.00	100.00 $\pm$ 0.00	100.00 $\pm$ 0.00
Negative control	30 ml	0.00 $\pm$ 0.00	0.00 $\pm$ 0.00	0.00 $\pm$ 0.00
CV (%)		4.3	3.67	3.24
P-value		<0.0001	<0.0001	<0.0001

**Note:** Means within a column followed by the same letter are not significantly different (Tukey Student Test (HSD) at  $p=0.05$ ).

**Abbreviations:** SE: Systematic Error; CV: Coefficient of variation; w/v: weight of powder/volume of water.

**Table 2:** Mean mortality  $\pm$  SE of *B. germanica* by botanical extracts and controls after 24, 48 and 72 hours exposure.

## CONCLUSION

Biopesticides are certain types of pesticides derived from animals, plants, bacteria, and certain minerals. They are natural products that are effective against bacteria, fungi, nematodes, viruses and insect pests. They are highly biodegradable, have varied modes of action, are no or less toxic to humans, are non-pollutant and they are readily available in the environment. They have various chemical properties and modes of action and effect on insects in different ways namely; repellents, feeding deterrents/anti-feedants, toxicants, growth retardants, chemosterilants, and attractants. So it is preferable to use the botanical insecticides instead of synthetic insecticides. So, we recommended using botanical insecticidal and being promoted and research is being conducted to find new sources of botanical insecticides.

## RECOMMENDATIONS

Continuous use of synthetic pesticides has resulted in negative effects such as pollution, health hazards and loss of biodiversity. For the reason that, botanical pesticides are successful in managing different pests with minimal costs and ecological side effects, easily available and biodegraded, have varied modes of action and low toxicity as well as availability of source materials. We, therefore, recommend using botanical insecticides as an integrated insect management program which can greatly reduce the use of synthetic insecticides.

## REFERENCES

- Pavela R. Larvicidal effects of various Euro-Asiatic plants against *Culex quinquefasciatus* say larvae (Diptera: Culicidae). *Parasitol Res.* 2010;102(3):555-559.
- Patil C, Patil S, Salunke B, Salunke R. Bioefficacy of *Plumbago zeylanica* (Plumbaginaceae) and *Cestrum nocturnum* (Solanaceae) plant extracts against *Aedes aegypti* (Diptera: Culicidae) and non-target fish *Poecilia reticulata*. *Parasitol Res.* 2011;108(5): 153-1263-2176.
- Ayodele M, Oke O. Studies on the potentials of some plant-based community pest management strategies in south west Nigeria. An investigation of the anti-termite potency of *Datura stramonium* L. *Int*

- J Agri Sci Sci Environ Technol (ASSET) Series B. 2003;2(2): 153-159.
4. Chowdhury N, Ghosh A, Chandra G. Mosquito larvicidal activities of *Solanum villosum* berry extract against the dengue vector *Stegomyia aegypti*. BMC Complement Altern Med. 2008;8:10.
  5. Hikal W, Baeshen R, Said H. Botanical insecticides as simple extractives for pest control. Cogent Biol. 2017;3(1):1404274.
  6. Ahmad W, Shilpa S, Sanjay K. Phytochemical screening and antimicrobial study of *Euphorbia hirta* extracts. J Med Plant Study. 2017;2:183-186.
  7. Joseph B, Sujatha S. Insight of botanical based biopesticides against economically important pest. Int J Pharmacol Life Sci. 2012;3(11): 2138-2148.
  8. Talukder F, Islam M, Hossain M, Rahman M, Alam M. Toxicity effects of botanicals and synthetic insecticides on *Tribolium castaneum* (Herbst) and *Rhyzopertha dominica* (Fabricius) Bangladesh. J Environ Sci. 2004;10(2):365-371.
  9. Rajashekar Y, Bakthavatsalam N, and Shivanandappa T. Botanicals as grain protectants. Psyche. 2012:1-13.
  10. Pavela R. Possibilities of botanical insecticide exploitation in plant protection. Pest Technol. 2007;1(4):47-52.
  11. Talukder F. Plant products as potential stored product insect management agents-A mini review. Emir J Food Agri. 2006;18(1): 17-32.
  12. Padin S, Fuse C, Urrutia M, DalBello G. Toxicity and repellency of nine medicinal plants against *Tribolium castaneum* in stored wheat. Bull Insectol. 2013;66(1):45-49.
  13. Koul O, Waliai S, Dhaliwal G. Essential oils as green pesticides: Potential and constraints. Biopestic Int. 2008;4(1):63-84.
  14. Navarro-Llopis V, Vacas S, Sanchis J, Primo J, Alfaro C. Chemosterilant bait stations coupled with sterile insect technique: An integrated strategy to control the Mediterranean fruit fly (Diptera: Tephritidae). J Econom Entomol. 2011;104(5):1647-1655.
  15. Wilke A, Nimmo D, John O, Kojin B, Capurro M, Marrelli M. Mini-review: Genetic enhancements to the sterile insect technique to control mosquito populations. As-Pac J Mol Biol Biotechnol. 2009;17(3):65-74.
  16. Yeom H, Kang S, Kim G, Park I. Insecticidal and acetylcholine esterase inhibition activity of Apiaceae plant essential oils and their constituents against adults of German cockroach (*Blattella germanica*). J Agri Food Chem. 2012;60(29):7194-7203.
  17. Slater J, James R, Pongracic J. Biological potency of German cockroach allergen extracts determined in an inner city population. Clin Exp Allergy. 2007;37(7):1033-1039.
  18. Mindykowski B. Cockroach allergens Per a 3 are oligomers. Dev Comp Immunol. 2010;34(7):722-733.
  19. Soedarto. Entomologi kedokteran. Penerbit buku kedokteran EGC. Jakarta. 1995.
  20. Brenner R, Koehler P, Patterson R. Health implications of cockroach infestations. Infect Med. 1987;4:349-358.
  21. Nasirian H. Rapid elimination of German cockroach, *Blattella germanica*, by fipronil and imidacloprid gel baits. J Arthropod Borne Dis. 2008;2(1):37-43.
  22. Alcamo I, Frishman A. The microbial flora of field collected cockroaches and other arthropods. J Environ Health. 1980;42:263-266.
  23. Lee CY, Yap HH, Chong NL. Insecticide toxicity on the adult German cockroach, *Blattella germanica* (L.) (Diptera: Blattellidae). Malaysian J Sci. 1996;17(1):1-9.
  24. Syed R, Manzoor, F, Adalat R, Abdul SA, Syed A. Laboratory evaluation of toxicity of insecticide formulations from different classes against American cockroach (Diptera: Blattidae). J Arthropod Borne Dis. 2014;8(1):21-34.
  25. Govindarajan M, Mathivanan T, Elumalai K, Krishnappa K, Anandan A. Mosquito larvicidal, ovicidal, and repellent properties of botanical extracts against *Anopheles stephensi*, *Aedes aegypti*, and *Culex quinquefasciatus* (Diptera: Culicidae). Parasitol Res. 2011;109:353-67.
  26. Amerasan D, Murugan K, Kovendan K. Adulticidal and repellent properties of indigenous plant extracts against *Culex quinquefasciatus* and *Aedes aegypti* (Diptera: Culicidae). Parasitol Res. 2012;110(5): 1607-1620.
  27. Muthukumar U, Govindarajan M, Rajeswary M, Hoti S. Synthesis and characterization of silver nanoparticles using *Gmelina asiatica* leaf extract against filariasis, dengue, and malaria vector mosquitoes. Parasitol Res. 2015;114(5):1817-1827.
  28. Pai H, Wu S, Hsu E. Insecticidal resistance in German cockroaches (*Blattella germanica*) from hospitals and households in Taiwan. Int J Environ Health Res. 2005;15(1):33-40.
  29. Kathy D. Pest management operators training manual. Classification, identification and biology of urban pests. 2016;1:9.
  30. Anis J, Primila K, Nisha V, Soory A, Sarika M. Safety of a *Indica* products to tetragynqathid spiders in rice ecosystem. J Biopestic. 2010;3:88-102.