

Controlling Nematodes for Sustainable Agriculture and Food Production

Bilton Ito*

Department of Entomology, University of Carbondale, Carbondale, USA

DESCRIPTION

The use and potential of Entomo Pathogenic Nematodes (EPNs) as a biological control agent in sustainable food production across a wide range of agricultural and other commodities is the focus of this research. Certain actions can be taken after the life history and behaviour of the pest are identified when contemplating prospective nematode management measures. Plant-parasitic nematodes are most abundant in the rooting zones of crops, typically in the top 30 cm of soil; nevertheless, worms can reach deeper soil layers, especially in unfavorable conditions, before returning to the surface layers when crop growth starts. Some species' free-living stages, such as the potato-cyst nematode, survive in unfavorable seasons through physiological or structural adaptation, or live for many seasons as eggs. When nematodes are actively hunting for host roots and surviving unfavorable growing seasons, they are normally at their most vulnerable. Free-living stages can survive longer without hosts than previously assumed, and unplanted periods of less than 12 months may be insufficient to manage species that reproduce quickly. Heat, desiccation, and anaerobes are the key abiotic variables that influence free-living nematodes and eggs. Plant-parasitic nematodes cannot survive in temperatures above 40°C for lengthy periods of time. During the summer season, these temperatures may be attained on bare soil in some regions, but heat penetration may not extend below 10 cm.

Flooding has been proposed as a method of nematode management in some areas. When the soil is drowned, all of the pore spaces are filled, restricting the oxygen supply to the soil micro flora and fauna. Many plant-parasitic nematodes are oxygen-intolerant and die quickly. When nematodes are held in deep water in the laboratory, similar effects are observed, but they may often be revived by aeration. In order to be effective in the field, the anaerobiosis must last long enough to kill the nematodes. Banana farms in Cote d'Ivoire are flooded for at least five weeks. Flooding is not an extensively utilized nematode management strategy, and it is unknown whether it is equally effective against all plant-parasitic species. The design of nematode management techniques necessitates a thorough understanding of the host plant's growth and the nematode's biology. In some climates, avoiding harm can be accomplished

by planting the crop at a time when nematode invasion is slow or when plant development is quicker than the parasite's. Seedlings produced in nematode-free nurseries or in soil-less potting media can reduce damage to transplanted short-cycle crops. In general, if a crop is protected from invasion during its early growth, it can generate a yield that is satisfactory to most farmers, even if it is planted in nematode-infested soil, as long as the crop is given appropriate water and nutrients. This will not prevent the growth of massive nematode populations on the root systems and in the soil, which will have an effect on the following crop. Nematodes (EPNs) can control pests thanks to a mutualistic relationship with bacteria that cause septicemia in the host and create a favorable environment for EPN development and reproduction. The diversity of EPNs found in Brazilian soils warrants additional investigation. Because of their mutualistic connection with bacteria of the genus *Xenorhabdus*, Entomo Pathogenic Nematodes (EPNs) of the families Steinernematidae and Heterorhabditidae can suppress pests. These nematodes enter the host through natural openings or the cuticle, carrying bacteria into the hemocele where they are capable to proliferate and kill the host within 24 to 48 hours and there by creating an environment conducive to nematode development and reproduction. When the insect host supplies run out, infective juveniles look for another host in the soil. Due to the inefficiency of traditional chemical and cultural strategies for insect soil management as well as the wide range of EPN hosts, interest in these biological control agents is growing. Entomo Pathogenic nematodes are now created in vivo and in vitro in several ways. Each method offers advantages and disadvantages in terms of manufacturing costs, technological know-how, economies of scale, and product quality, and each method has the potential to be improved. Following manufacture, there are a range of formulation and application options. The goals of this review are to give a summary and analysis of elements that influence EPN production and application, as well as insight into how to improve techniques to achieve more biological control success. One should sanitize the crop regularly in order to prevent the nematodes. We can also reduce it by using some methods like crop rotation, fallowing, and soil solarization. But these are effective only for 11-12 months. Nematicides are commonly used pesticides to kill nematodes.

Correspondence to: Bilton Ito, Department of Entomology, University of Carbondale, Carbondale, USA, E-mail: bילו786@gmail.com

Received: December 03, 2021; **Accepted:** December 17, 2021; **Published:** December 24, 2021

Citation: Ito B (2021) Controlling Nematodes for Sustainable Agriculture and Food Production. Entomol Ornithol Herpetol. 10:264.

Copyright: © 2021 Ito B. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.